

International Union of Pharmacology. XXII. Nomenclature for Chemokine Receptors

PHILIP M. MURPHY,¹ MARCO BAGGIOLINI, ISRAEL F. CHARO, CAROLINE A. HEBERT, RICHARD HORUK,
 KOUJI MATSUSHIMA, LOUIS H. MILLER, JOOST J. OPPENHEIM, AND CHRISTINE A. POWER

Laboratory of Host Defenses, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland (P.M.M.); Theodor-Kocher Institute, Bern, Switzerland (M.B.); Gladstone Institute of Cardiovascular Disease, San Francisco, California (I.F.C.); Genentech, Inc., South San Francisco, California (C.A.H.); Department of Immunology, Berlex Biosciences, Richmond, California (R.H.); Department of Molecular Preventive Medicine, School of Medicine, University of Tokyo, Bunkyo, Tokyo, Japan (K.M.); Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland (L.H.M.); Laboratory of Molecular Immunoregulation, Division of Basic Science, National Cancer Institute-Frederick Cancer Research and Development Center, Frederick, Maryland (J.J.O.); and Sero Pharmaceutical Research Institute, Plan-les-Ouates, Geneva, Switzerland (C.A.P.)

This paper is available online at <http://www.pharmrev.org>

Abstract	146
I. Overview	146
II. Introduction	147
A. Historical background	147
B. Chemokine classification	147
C. Chemokine receptor classification and nomenclature	151
D. Chemokine receptor structure	151
E. Chemokine receptor specificity for ligands and leukocytes	153
III. CXCR chemokine receptor subtypes	155
A. CXCR1 and CXCR2	155
B. CXCR3	157
C. CXCR4	157
D. CXCR5	159
IV. CC chemokine receptor subtypes	159
A. CCR1	159
B. CCR2	160
C. CCR3	161
D. CCR4	162
E. CCR5	162
F. CCR6	164
G. CCR7	164
H. CCR8	165
I. CCR9	165
J. CCR10	165
K. CCR11	165
V. CX3CR chemokine receptor subtypes	166
A. CX3CR1	166
VI. C chemokine receptor subtypes	166
A. XCR1	166
VII. Chemokine binding proteins	166
A. Duffy	166
B. D6	167
VIII. Virus-encoded chemokine receptors	167
A. ECRF3	167
B. US28	167
C. KSHV GPCR	168

¹ Address for correspondence: Dr. Philip M. Murphy, Laboratory of Host Defenses, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Building 10, Room 11N113, Bethesda, MD 20892. E-mail: pmm@nih.gov

D. UL12	168
E. E1	168
IX. Conclusions	168
Acknowledgments	168
References	168

Abstract—Chemokine receptors comprise a large family of seven transmembrane domain G protein-coupled receptors differentially expressed in diverse cell types. Biological activities have been most clearly defined in leukocytes, where chemokines coordinate development, differentiation, anatomic distribution, trafficking, and effector functions and thereby regulate innate and adaptive immune responses. Pharmacological analysis of chemokine receptors is at an early stage of development. Disease indications have been established in human immunodeficiency virus/acquired immune deficiency syndrome and in *Plasmodium vivax* malaria, due to exploitation of CCR5 and Duffy, respectively, by the pathogen for cell entry. Additional indications are emerging among inflammatory and immunologically mediated diseases, but selection of targets in this area still remains somewhat speculative. Small molecule antagonists with nanomole-

lar affinity have been reported for 7 of the 18 known chemokine receptors but have not yet been studied in clinical trials. Virally encoded chemokine receptors, as well as chemokine agonists and antagonists, and chemokine scavengers have been identified in medically important poxviruses and herpesviruses, again underscoring the importance of the chemokine system in microbial pathogenesis and possibly identifying specific strategies for modulating chemokine action therapeutically. The purpose of this review is to update current concepts of the biology and pharmacology of the chemokine system, to summarize key information about each chemokine receptor, and to describe a widely accepted receptor nomenclature system, ratified by the International Union of Pharmacology, that is facilitating clear communication in this area.

I. Overview

The aim of this article is to describe the nomenclature system for chemokine receptors, as approved by the Nomenclature Committee of the International Union of Pharmacology (NC-IUPHAR),² and to update their main molecular and biological properties. A general overview is given first, followed by a synopsis of key information about each receptor, with an emphasis on recent discoveries and new concepts.

Chemokine receptors are defined by their ability to signal on binding one or more members of the chemokine superfamily of chemotactic cytokines (Premack and Schall, 1996; Baggiolini et al., 1997; Yoshie et al., 1997; Luster, 1998; Zlotnik et al., 1999). To date, 18 human proteins have met this definition, and they have been designated CXCR1 through 5, CCR1 through 11, XCR1, and CX3CR1 based on their specific chemokine preferences, as described in subsequent sections. Together, chemokine receptors comprise a large branch of the rhodopsin family of cell surface, seven-transmembrane domain (7TMD), G protein-coupled receptors (GPCRs). In addition, D6 and Duffy (sometimes called the Duffy antigen receptor for chemokines, or DARC) are 7TMD

mammalian chemokine-binding proteins that apparently do not signal and therefore are excluded from the systematic nomenclature (Horuk, 1994; Nibbs et al., 1997a).

To date, chemokine receptor-like sequences have been identified in mammals, birds (Gupta et al., 1998a), and fish (Daniels et al., 1999) but not in invertebrates, plants, yeast, or bacteria, suggesting a relatively recent origin. Common features include conserved structure [25–80% amino acid (aa) identity], coupling to the G_i class of G proteins, expression in leukocytes, and chemotactic signaling. The major shared biological function is leukocyte trafficking and dependent processes such as immune surveillance, innate and adaptive immune responses, and various forms of pathological inflammation (Springer, 1994; Foxman et al., 1997). Within this general area, however, each chemokine receptor appears to have a specific role, determined by its expression pattern on specific subsets of leukocytes, and by the temporal and spatial specificity of cognate ligand expression. Specific roles have also been delineated in hematopoiesis (Broxmeyer et al., 1996, 1999; Reid et al., 1999), angiogenesis (Salcedo et al., 1999), development (Forster et al., 1996; Nagasawa et al., 1996; Ma et al., 1998; Tachibana et al., 1998; Zou et al., 1998), and, counterintuitively, facilitation of certain infectious diseases.

With regard to the latter, two major themes have been defined. In the first, cellular chemokine receptors are exploited as cell entry and disease transmission factors by intracellular pathogens. Rigorously proved examples of this are the human immunodeficiency virus (HIV)

² Abbreviations: NC-IUPHAR, Nomenclature Committee of the International Union of Pharmacology; 7TMD, seven-transmembrane domain; HHV8, human herpesvirus 8; KS, Kaposi's sarcoma; NK, natural killer; GPCR, G protein-coupled receptor; HIV, human immunodeficiency virus; AIDS, acquired immune deficiency syndrome; mAb, monoclonal antibody; IL, interleukin; ORF, open reading frame; AOP, amino-oxypentane; aa, amino acid(s); HCMV, human cytomegalovirus.

coreceptor CCR5 in acquired immune deficiency syndrome (AIDS) and Duffy in the form of malaria caused by *Plasmodium vivax*; CXCR4 and other chemokine receptors also function as HIV coreceptors, but their importance in disease is not established (Horuk et al., 1994; Rucker et al., 1997; Berger et al., 1999). The second theme, which is not as well understood, involves herpesvirus- and poxvirus-encoded chemokines and chemokine receptors, evidently acquired as copied host genes, which may subvert the immune response or dysregulate cell growth (reviewed in Pease and Murphy, 1998).

Apart from Duffy in malaria and CCR5 in HIV/AIDS, disease indications have not yet been unequivocally established for chemokine receptors. Rapid progress in this area can be anticipated in the near future as receptor knockout mice and receptor-blocking agents are tested in animal models of disease. To date, only CXCR4 has been shown to be essential for life. Phenotypes of knockout mice for other chemokine receptors are more subtle in the absence of specific stresses (Gerard, 1999). Many types of chemokine and chemokine receptor-blocking agents of high and low selectivity have been discovered, including viral chemokine scavengers, viral chemokine antagonists, antagonistic chemokine variants, small molecules, ribozymes, intrakinases, and monoclonal antibodies (mAbs) (Schwarz and Wells, 1999). Moreover, a novel HIV vaccine has been discovered in which CCR5 is a critical component (Lacasse et al., 1999). However, as of this writing, none of these has been tested in a clinical trial.

II. Introduction

A. Historical Background

To understand chemokine receptors, first their ligands must be explained. Chemokines are perhaps the most complex of GPCR ligands because of their large number, overlapping receptor specificity, and extensive phylogenetic divergence. To date, more than 40 different human chemokines have been identified, with the first identified in 1977, when Walz et al. (1977) sequenced native platelet factor 4, a procoagulant and angiostatic factor stored in platelet α -granules. Subsequently, from 1984 through 1989, cDNAs for structurally related proteins, including IP-10 (see Table 1 for chemokine acronyms; synonyms and chemokine classes are given in Table 2), JE, Mig, RANTES (regulated on activation, normal T cell expressed and secreted), I-309, KC, and macrophage inflammatory protein-1 α (MIP-1 α), were cloned by investigators looking mainly for cell differentiation- and activation-associated genes, establishing the existence of a gene family before identifying any functions (Wolpe and Cerami, 1989; Schall, 1991; Oppenheim et al., 1991).

The discovery of the neutrophil-targeted chemokine interleukin (IL)-8 represents a landmark in immunology

because it was the first leukocyte subtype-selective chemoattractant to be found (Yoshimura et al., 1987; Walz et al., 1987). The discovery of IL-8 also focused the search for functions for other chemokines on leukocyte chemotaxis and stimulated a search for new family members. Interest in the field grew with subsequent reports of MCP-1, RANTES, and eotaxin, the first important monocyte-, T cell-, and eosinophil-directed chemokines, respectively (Matsumura et al., 1989; Yoshimura et al., 1989; Schall et al., 1990; Jose et al., 1994). Methods of chemokine discovery expanded to include purification of chemoattractant activity as well as cDNA cloning by signal sequence trapping, homology hybridization, and, most recently, bioinformatics and expressed sequence tag (EST) analysis (Tashiro et al., 1993, 1999; Wells and Peitsch, 1997). Chemokines are particularly easy to find in EST databases because their coding sequences are sufficiently small, typically 70 to 90 codons, to be captured by a single EST and because their conserved sequence motifs are easy to recognize (see later). As the number of family members expanded, various short-lived collective terms for them were used, including "the platelet factor-4 family" (Wolpe and Cerami, 1989), "the small inducible cytokine family" (Schall, 1991), and "the intercrines" (Oppenheim et al., 1991). Finally, in 1992 at the Third International Symposium on Chemotactic Cytokines in Baden, the term "chemokine," a neologism short for "chemotactic cytokines," was coined and accepted as the standard (Lindley et al., 1993).

With respect to leukocyte specificity, both broad- and narrow-spectrum chemokines have been identified. Together they cover the full spectrum of leukocytes, acting through a signaling pathway that includes a pertussis toxin-sensitive G protein (G_i/G_o), calcium flux, and chemotaxis. This fact pointed to use of GPCRs and suggested homology hybridization as a strategy to identify them (reviewed in Murphy, 1996), which has been highly successful.

In 1995, an NC-IUPHAR subcommittee on chemokine receptor nomenclature was organized. Recommendations developed at the second Gordon Conference on Chemotactic Cytokines held in Plymouth, NH, in 1996, were accepted unanimously by meeting participants, ratified by NC-IUPHAR in January 1997, and widely used since. The nomenclature is based on the subclassification of the chemokine superfamily, delineated in the next section. In 1998, a second nomenclature committee, led by O. Yoshie and A. Zlotnik, was formed to address the proliferation of chemokine aliases that has accompanied the codiscovery of chemokines by multiple groups using bioinformatics (Table 1). A nomenclature system that parallels the receptor nomenclature was proposed at the Keystone Symposium on Chemokine and Chemokine Receptors, January 18 to 23, 1999, in Keystone, CO (Table 2).

TABLE 1

Chemokine acronyms

Synonyms and chemokine class are given in Table 2.

6CKine, chemokine with 6 cysteines
AMAC, alternative macrophage activation-associated CC chemokine
ATAC, activation-induced, chemokine-related molecule exclusively expressed in CD8 ⁺ T lymphocytes
BCA-1, B cell-activating chemokine-1
BLC, B lymphocyte chemoattractant
BRAC, breast and kidney chemokine
CC-#, CC chemokine-#
COCK-#, CC chemokine-#
ckβ#, CC chemokine #
CRG, cytokine-responsive gene
CTAP III, connective tissue-activating peptide III
deβ-ck-1, dendritic cell β-chemokine-1
DC-CK-1, dendritic cell chemokine 1
ELC, Epstein-Barr virus-induced receptor ligand chemokine
ENA-78, epithelial cell-derived neutrophil-activating factor, 78 amino acids
FIC, fibroblast-inducible cytokine
GCP-#, granulocyte chemoattractant protein-#
Gro, growth-related oncoene
HCC-#, hemofiltrate CC chemokine-#
IL-8, interleukin-8
γIP-10, γ-interferon-inducible protein-10
I-TAC, interferon-inducible T cell α-chemoattractant
LARC, liver- and activation-related chemokine
LCC, liver CC chemokine
LEC, liver-expressed chemokine
Lkn-1, leukotactin-1
LMC, lymphocyte and monocyte chemoattractant CC chemokine
LYNAP, lymphocyte-derived neutrophil-activating peptide
MARC, mast cell activation-related chemokine
MCAF, monocyte chemoattractant and activating factor
MCIF, macrophage colony-inhibitory factor
MCP-#, monocyte chemoattractant protein-#
MDC, macrophage-derived chemokine
MDNCF, monocyte-derived neutrophil chemotactic factor
MGSA, melanoma growth-stimulatory activity
Mig, monokine induced by γ-interferon
MIP-#, macrophage inflammatory protein-#
MPiF-#, myeloid progenitor inhibitory factor-#
MRP-#, MIP-related protein-#
Mtn-1, monotactin-1
NAF, neutrophil-activating factor
NAP-#, neutrophil-activating protein-#
NCC-#, novel CC chemokine-#
PARC, pulmonary- and activation-regulated chemokine
PBP, platelet basic protein
PBSF, pre-B cell-stimulatory factor
PF-4, platelet factor-4
RANTES, regulated on activation normal T cell expressed and secreted
SCM-1, single C motif-1
SCY#, small cytokine #
SDF-1, stromal cell-derived factor-1
SIS-#, small inducible secreted protein-#
SLC, secondary lymphoid tissue chemokine
STCP-1, stimulated T cell chemoattractant protein-1
TARC, thymus- and activation-related chemokine
TCA-#, T-cell activation protein-#
TECK, thymus-expressed chemokine
TG, thromboglobulin
TLSF, thymic lymphoma cell-stimulating factor
TPAR, TPA-repressed protein

B. Chemokine Classification

Chemokines can be subclassified by structure according to the number and spacing of conserved cysteines into four major groups, given the preferred names CXC, CC, C, and CX3C, which are used in the systematic nomenclatures (Tables 2 and 3). Less commonly these groups are referred to by the Greek letters α , β , γ , and δ , respectively. CXC, CC, and CX3C chemokines all have four conserved cysteines, whereas C chemokines have only two, corresponding to the second and fourth cysteines in the other groups (Fig. 1). A small subgroup of CC chemokines has six cysteines. CXC and CX3C chemokines are distinguished by the presence of one (CXC) or three (CX3C) aa between the first and second cysteines, whereas the first two cysteines of CC chemokines are adjacent. Both the CC and CXC groups have many known members, whereas human lymphotactin α and β (Kelner et al., 1994) and fractalkine (Bazan et al., 1997) and their equivalents in other species are the only known examples of C and CX3C chemokines, respectively. A cDNA encoding a CXC chemokine-like protein has been discovered in lamprey, suggesting that the origin of the family, and possibly the division into subclasses, is ancient (Najakshin et al., 1999).

Fractalkine is an interesting model for how chemokines may be presented to target cells. It has a multimolecular structure consisting of a chemokine domain fused to a mucin-like stalk plus a transmembrane domain, which anchors the molecule to the plasma membrane, and a cytoplasmic domain. Consistent with this, it functions as an adhesion molecule by binding directly to CX3CR1 (Imai et al., 1997b). Fractalkine also induces cell migration as either a tethered or shed ligand. Although other chemokines lack a transmembrane domain and are secreted, they are able to use glycosaminoglycans for tethering to plasma membrane. This provides a mechanism for gradient formation under conditions of high blood flow. Once "posted" in this manner, chemokines may be "read" by passing leukocytes, which then activate β_2 -integrins, bind to endothelium and transmigrate from blood to tissue (Tanaka et al., 1993). However, fractalkine is the only chemokine shown to act as a direct cell adhesion molecule.

CXC chemokines can be further subclassified by structure into ELR+ and ELR- molecules based on the presence or absence of the tripeptide motif glutamic acid-leucine-arginine (ELR) N-terminal to the first cysteine. This provides the only strong functional correlate of the structural classification: the specificity of ELR+ CXC chemokines for neutrophils (Hebert et al., 1991). A second classification scheme based on function and expression pattern has also been proposed. It includes an inflammatory/inducible group, which is regulated by proinflammatory stimuli such as lipopolysaccharide and primary cytokines such as IL-1 and tumor necrosis factor, and which together orchestrate innate and adaptive

[illegible]

Official Name	Chromosome	Protein	Gene	Common Synonyms	Other Names	Disease Association	Subclass
						Number	
						Human	
CXCL1 (Chemokine)	4q21		GRO1	GRO α , MGSX, (monocyte) NS1/KC α , (monocyte) MIP-2 α	SCYB1; NAP-3; GRO1 monocyte	P13650P10869	ELR+; ELR+; ELR+
CXCL2	4q21		GRO2	GRO β , MIP-2 α	SCYB2; GRO2 monocyte	P19841	ELR+
CXCL3	4q21		GRO3	GRO γ , MIP-2 α	SCYB3; (GRO3 monocyte)	P19876	ELR+
CXCL4	4q11-q23		GRO4	GRO δ , MIP-2 α	SCYB4; (GRO4 monocyte)	P42630	ELR+
CXCL5	4q11-q21		SCYB5	ENA-78	SCYB5	P490182	ELR+
CXCL6	4q12-q13		SCYB6	GCP-2	SCYB6	P27775	ELR+
CXCL7	4q12-q13		PP3P	PPP-2/CXAP-III β -PTG-SNAP-2 β	SCYB7; low-affinity platelet factor-4	P27775	ELR+
CXCL8	4q12-q13		SCYB8	MIP-3 α , MIP-3 β	SCYB8 (monocyte) GRO-1 β	P27775	ELR+
CXCL9	4q21		ARF	ARF	SCYB9 (monocyte) GRO-1 α	P27775	ELR+
CXCL10	4q21		INP10	IFP-1 β , (monocyte) CCR-2	SCYB10; (monocyte) CCR-2	P10696	ELR-
CXCL11	4q21.2		IFP1	IFPAC, pART, IFP β , H174	SCYB11	P10696	ELR-
CXCL12	4q21.2		IFP2	IFPAC, pART, IFP β , IFP2	SCYB11	P10696	ELR-
CXCL13	4q21.2		IFP3	IFPAC, pART, IFP β , IFP3	SCYB13	P10696	ELR-
CXCL14	4q21.2		IFP4	IFPAC, pART, IFP β , IFP4	SCYB14	P10696	ELR-
CXCL15	4q21.2		IFP5	IFPAC, pART, IFP β , IFP5	SCYB15	P10696	ELR-
CXCL16	4q21.2		IFP6	IFPAC, pART, IFP β , IFP6	SCYB16	P10696	ELR-
CXCL17	4q21.2		IFP7	IFPAC, pART, IFP β , IFP7	SCYB17	P10696	ELR-
CXCL18	4q21.2		IFP8	IFPAC, pART, IFP β , IFP8	SCYB18	P10696	ELR-
CXCL19	4q21.2		IFP9	IFPAC, pART, IFP β , IFP9	SCYB19	P10696	ELR-
CXCL20	4q21.2		IFP10	IFPAC, pART, IFP β , IFP10	SCYB20	P10696	ELR-
CXCL21	4q21.2		IFP11	IFPAC, pART, IFP β , IFP11	SCYB21	P10696	ELR-
CXCL22	4q21.2		IFP12	IFPAC, pART, IFP β , IFP12	SCYB22	P10696	ELR-
CXCL23	4q21.2		IFP13	IFPAC, pART, IFP β , IFP13	SCYB23	P10696	ELR-
CXCL24	4q21.2		IFP14	IFPAC, pART, IFP β , IFP14	SCYB24	P10696	ELR-
CXCL25	4q21.2		IFP15	IFPAC, pART, IFP β , IFP15	SCYB25	P10696	ELR-
CXCL26	4q21.2		IFP16	IFPAC, pART, IFP β , IFP16	SCYB26	P10696	ELR-
CXCL27	4q21.2		IFP17	IFPAC, pART, IFP β , IFP17	SCYB27	P10696	ELR-
CXCL28	4q21.2		IFP18	IFPAC, pART, IFP β , IFP18	SCYB28	P10696	ELR-
CXCL29	4q21.2		IFP19	IFPAC, pART, IFP β , IFP19	SCYB29	P10696	ELR-
CXCL30	4q21.2		IFP20	IFPAC, pART, IFP β , IFP20	SCYB30	P10696	ELR-
CXCL31	4q21.2		IFP21	IFPAC, pART, IFP β , IFP21	SCYB31	P10696	ELR-
CXCL32	4q21.2		IFP22	IFPAC, pART, IFP β , IFP22	SCYB32	P10696	ELR-
CXCL33	4q21.2		IFP23	IFPAC, pART, IFP β , IFP23	SCYB33	P10696	ELR-
CXCL34	4q21.2		IFP24	IFPAC, pART, IFP β , IFP24	SCYB34	P10696	ELR-
CXCL35	4q21.2		IFP25	IFPAC, pART, IFP β , IFP25	SCYB35	P10696	ELR-
CXCL36	4q21.2		IFP26	IFPAC, pART, IFP β , IFP26	SCYB36	P10696	ELR-
CXCL37	4q21.2		IFP27	IFPAC, pART, IFP β , IFP27	SCYB37	P10696	ELR-
CXCL38	4q21.2		IFP28	IFPAC, pART, IFP β , IFP28	SCYB38	P10696	ELR-
CXCL39	4q21.2		IFP29	IFPAC, pART, IFP β , IFP29	SCYB39	P10696	ELR-
CXCL40	4q21.2		IFP30	IFPAC, pART, IFP β , IFP30	SCYB40	P10696	ELR-
CXCL41	4q21.2		IFP31	IFPAC, pART, IFP β , IFP31	SCYB41	P10696	ELR-
CXCL42	4q21.2		IFP32	IFPAC, pART, IFP β , IFP32	SCYB42	P10696	ELR-
CXCL43	4q21.2		IFP33	IFPAC, pART, IFP β , IFP33	SCYB43	P10696	ELR-
CXCL44	4q21.2		IFP34	IFPAC, pART, IFP β , IFP34	SCYB44	P10696	ELR-
CXCL45	4q21.2		IFP35	IFPAC, pART, IFP β , IFP35	SCYB45	P10696	ELR-
CXCL46	4q21.2		IFP36	IFPAC, pART, IFP β , IFP36	SCYB46	P10696	ELR-
CXCL47	4q21.2		IFP37	IFPAC, pART, IFP β , IFP37	SCYB47	P10696	ELR-
CXCL48	4q21.2		IFP38	IFPAC, pART, IFP β , IFP38	SCYB48	P10696	ELR-
CXCL49	4q21.2		IFP39	IFPAC, pART, IFP β , IFP39	SCYB49	P10696	ELR-
CXCL50	4q21.2		IFP40	IFPAC, pART, IFP β , IFP40	SCYB50	P10696	ELR-
CXCL51	4q21.2		IFP41	IFPAC, pART, IFP β , IFP41	SCYB51	P10696	ELR-
CXCL52	4q21.2		IFP42	IFPAC, pART, IFP β , IFP42	SCYB52	P10696	ELR-
CXCL53	4q21.2		IFP43	IFPAC, pART, IFP β , IFP43	SCYB53	P10696	ELR-
CXCL54	4q21.2		IFP44	IFPAC, pART, IFP β , IFP44	SCYB54	P10696	ELR-
CXCL55	4q21.2		IFP45	IFPAC, pART, IFP β , IFP45	SCYB55	P10696	ELR-
CXCL56	4q21.2		IFP46	IFPAC, pART, IFP β , IFP46	SCYB56	P10696	ELR-
CXCL57	4q21.2		IFP47	IFPAC, pART, IFP β , IFP47	SCYB57	P10696	ELR-
CXCL58	4q21.2		IFP48	IFPAC, pART, IFP β , IFP48	SCYB58	P10696	ELR-
CXCL59	4q21.2		IFP49	IFPAC, pART, IFP β , IFP49	SCYB59	P10696	ELR-
CXCL60	4q21.2		IFP50	IFPAC, pART, IFP β , IFP50	SCYB60	P10696	ELR-
CXCL61	4q21.2		IFP51	IFPAC, pART, IFP β , IFP51	SCYB61	P10696	ELR-
CXCL62	4q21.2		IFP52	IFPAC, pART, IFP β , IFP52	SCYB62	P10696	ELR-
CXCL63	4q21.2		IFP53	IFPAC, pART, IFP β , IFP53	SCYB63	P10696	ELR-
CXCL64	4q21.2		IFP54	IFPAC, pART, IFP β , IFP54	SCYB64	P10696	ELR-
CXCL65	4q21.2		IFP55	IFPAC, pART, IFP β , IFP55	SCYB65	P10696	ELR-
CXCL66	4q21.2		IFP56	IFPAC, pART, IFP β , IFP56	SCYB66	P10696	ELR-
CXCL67	4q21.2		IFP57	IFPAC, pART, IFP β , IFP57	SCYB67	P10696	ELR-
CXCL68	4q21.2		IFP58	IFPAC, pART, IFP β , IFP58	SCYB68	P10696	ELR-
CXCL69	4q21.2		IFP59	IFPAC, pART, IFP β , IFP59	SCYB69	P10696	ELR-
CXCL70	4q21.2		IFP60	IFPAC, pART, IFP β , IFP60	SCYB70	P10696	ELR-
CXCL71	4q21.2		IFP61	IFPAC, pART, IFP β , IFP61	SCYB71	P10696	ELR-
CXCL72	4q21.2		IFP62	IFPAC, pART, IFP β , IFP62	SCYB72	P10696	ELR-
CXCL73	4q21.2		IFP63	IFPAC, pART, IFP β , IFP63	SCYB73	P10696	ELR-
CXCL74	4q21.2		IFP64	IFPAC, pART, IFP β , IFP64	SCYB74	P10696	ELR-
CXCL75	4q21.2		IFP65	IFPAC, pART, IFP β , IFP65	SCYB75	P10696	ELR-
CXCL76	4q21.2		IFP66	IFPAC, pART, IFP β , IFP66	SCYB76	P10696	ELR-
CXCL77	4q21.2		IFP67	IFPAC, pART, IFP β , IFP67	SCYB77	P10696	ELR-
CXCL78	4q21.2		IFP68	IFPAC, pART, IFP β , IFP68	SCYB78	P10696	ELR-
CXCL79	4q21.2		IFP69	IFPAC, pART, IFP β , IFP69	SCYB79	P10696	ELR-
CXCL80	4q21.2		IFP70	IFPAC, pART, IFP β , IFP70	SCYB80	P10696	ELR-
CXCL81	4q21.2		IFP71	IFPAC, pART, IFP β , IFP71	SCYB81	P10696	ELR-
CXCL82	4q21.2		IFP72	IFPAC, pART, IFP β , IFP72	SCYB82	P10696	ELR-
CXCL83	4q21.2		IFP73	IFPAC, pART, IFP β , IFP73	SCYB83	P10696	ELR-
CXCL84	4q21.2		IFP74	IFPAC, pART, IFP β , IFP74	SCYB84	P10696	ELR-
CXCL85	4q21.2		IFP75	IFPAC, pART, IFP β , IFP75	SCYB85	P10696	ELR-
CXCL86	4q21.2		IFP76	IFPAC, pART, IFP β , IFP76	SCYB86	P10696	ELR-
CXCL87	4q21.2		IFP77	IFPAC, pART, IFP β , IFP77	SCYB87	P10696	ELR-
CXCL88	4q21.2		IFP78	IFPAC, pART, IFP β , IFP78	SCYB88	P10696	ELR-
CXCL89	4q21.2		IFP79	IFPAC, pART, IFP β , IFP79	SCYB89	P10696	ELR-
CXCL90	4q21.2		IFP80	IFPAC, pART, IFP β , IFP80	SCYB90	P10696	ELR-
CXCL91	4q21.2		IFP81	IFPAC, pART, IFP β , IFP81	SCYB91	P10696	ELR-
CXCL92	4q21.2		IFP82	IFPAC, pART, IFP β , IFP82	SCYB92	P10696	ELR-
CXCL93	4q21.2		IFP83	IFPAC, pART, IFP β , IFP83	SCYB93	P10696	ELR-
CXCL94	4q21.2		IFP84	IFPAC, pART, IFP β , IFP84	SCYB94	P10696	ELR-
CXCL95	4q21.2		IFP85	IFPAC, pART, IFP β , IFP85	SCYB95	P10696	ELR-
CXCL96	4q21.2		IFP86	IFPAC, pART, IFP β , IFP86	SCYB96	P10696	ELR-
CXCL97	4q21.2		IFP87	IFPAC, pART, IFP β , IFP87	SCYB97	P10696	ELR-
CXCL98	4q21.2		IFP88	IFPAC, pART, IFP β , IFP88	SCYB98	P10696	ELR-
CXCL99	4q21.2		IFP89	IFPAC, pART, IFP β , IFP89	SCYB99	P10696	ELR-
CXCL100	4q21.2		IFP90	IFPAC, pART, IFP β , IFP90	SCYB100	P10696	ELR-
CXCL101	4q21.2		IFP91	IFPAC, pART, IFP β , IFP91	SCYB101	P10696	ELR-
CXCL102	4q21.2		IFP92	IFPAC, pART, IFP β , IFP92	SCYB102	P10696	ELR-
CXCL103	4q21.2		IFP93	IFPAC, pART, IFP β , IFP93	SCYB103	P10696	ELR-
CXCL104	4q21.2		IFP94	IFPAC, pART, IFP β , IFP94	SCYB104	P10696	ELR-
CXCL105	4q21.2		IFP95	IFPAC, pART, IFP β , IFP95	SCYB105	P10696	ELR-
CXCL106	4q21.2		IFP96	IFPAC, pART, IFP β , IFP96	SCYB106	P10696	ELR-
CXCL107	4q21.2		IFP97	IFPAC, pART, IFP β , IFP97	SCYB107	P10696	ELR-
CXCL108	4q21.2		IFP98	IFPAC, pART, IFP β , IFP98	SCYB108	P10696	ELR-
CXCL109	4q21.2		IFP99	IFPAC, pART, IFP β , IFP99	SCYB109	P10696	ELR-
CXCL110	4q21.2		IFP100	IFPAC, pART, IFP β , IFP100	SCYB110	P10696	ELR-
CXCL111	4q21.2		IFP101	IFPAC, pART, IFP β , IFP101	SCYB111	P10696	ELR-
CXCL112	4q21.2		IFP102	IFPAC, pART, IFP β , IFP102	SCYB112	P10696	ELR-
CXCL113	4q21.2		IFP103	IFPAC, pART, IFP β , IFP103	SCYB113	P10696	ELR-
CXCL114	4q21.2		IFP104	IFPAC, pART, IFP β , IFP104	SCYB114	P10696	ELR-
CXCL115	4q21.2		IFP105	IFPAC, pART, IFP β , IFP105	SCYB115	P10696	ELR-
CXCL116	4q21.2		IFP106	IFPAC, pART, IFP β , IFP106	SCYB116	P10696	ELR-
CXCL117	4q21.2		IFP107	IFPAC, pART, IFP β , IFP107	SCYB117	P10696	ELR-
CXCL118	4q21.2		IFP108	IFPAC, pART, IFP β , IFP108	SCYB118	P10696	ELR-
CXCL119	4q21.2		IFP109	IFPAC, pART, IFP β , IFP109	SCYB119	P10696	ELR-
CXCL120	4q21.2		IFP110	IFPAC, pART, IFP β , IFP110	SCYB120	P10696	ELR-
CXCL121	4q21.2		IFP111	IFPAC, pART, IFP β , IFP111	SCYB121	P10696	ELR-
CXCL122	4q21.2		IFP112	IFPAC, pART, IFP β , IFP112	SCYB122	P10696	ELR-
CXCL123	4q21.2		IFP113	IFPAC, pART, IFP β , IFP113	SCYB123	P10696	ELR-
CXCL124	4q21.2		IFP114	IFPAC, pART, IFP β , IFP114	SCYB124	P10696	ELR-
CXCL125	4q21.2		IFP115	IFPAC, pART, IFP β , IFP115	SCYB125	P10696	ELR-
CXCL126	4q21.2		IFP116	IFPAC, pART, IFP β , IFP116	SCYB126	P10696	ELR-
CXCL127	4q21.2		IFP117	IFPAC, pART, IFP β , IFP117	SCYB127	P10696	ELR-
CXCL128	4q21.2		IFP118	IFPAC, pART, IFP β , IFP118	SCYB128	P10696	ELR-
CXCL129	4q21.2		IFP119	IFPAC, pART, IFP β , IFP119	SCYB129	P10696	ELR-
CXCL130	4q21.2		IFP120	IFPAC, pART, IFP β , IFP120	SCYB130	P10696	ELR-
CXCL131	4q21.2		IFP121	IFPAC, pART, IFP β , IFP121	SCYB131	P10696	ELR-
CXCL132	4q21.2		IFP122	IFPAC, pART, IFP β , IFP122	SCYB132	P10696	ELR-
CXCL133	4q21.2		IFP123	IFPAC, pART, IFP β , IFP123	SCYB133	P10696	ELR-
CXCL134	4q21.2		IFP124	IFPAC, pART, IFP β , IFP124	SCYB134	P10696	ELR-
CXCL135	4q21.2		IFP125	IFPAC, pART, IFP β , IFP125	SCYB135	P10696	ELR-
CXCL136	4q21.2		IFP126	IFPAC, pART, IFP β , IFP126	SCYB136	P10696	ELR-

^a KC and MIP-2 are mouse proteins with similar sequence relatedness to each of the three human Gro proteins. Sequential N-terminal truncation of PPP produces the chemokines shown. Only NAP-2 has leukocyte chemoattractant activity, specifically for neutrophils.

Sequential N-terminal truncation of FGF products are shown in Table 1. Only NAT-2 has neurotrophic activity, indicating that the neurotrophic activity of NAT-1 is due to the presence of the C-terminal sequence. The amino acid sequence of NAT-2 is identical to the sequence of the C-terminal region of the mature FGF-10 protein (100–165) [10].

^d SDF-1 α and SDF-1 β are splice variants of the same human gene but appear to be functionally equivalent.

TABLE 3
Chemokine receptors; nomenclature, pharmacology and biology

N.D., not determined; N.A., not available. Two splice variants affecting the ORF have been identified for each of the following receptors: CXCR4, CXCR5, CXCR2, and CXCR9; however, biological or pharmacological significance has not been determined. Other HIV coreceptors include US28 (Pleschke et al., 1997), STRL331 (Bonzio/TYMSTR (Deng et al., 1997; Parzan et al., 1997), ROC1 (Shimizu et al., 2000); and ChemR23 (Samson et al., 1998).

Name	Previous Names	Gene Chromosome	Accession Number		Pharmacology		Biology	
			Human	Mouse	Selective Chemokine	Selective Neutrophils (agonist), Macrophages (antagonist)	Major Phenotypes of Chemically Deficient Mice/Humans	HIV Coreceptor Activity
CXCR1 chemokine receptors	IL8R _α , IL-8R ₁ , IL-8R _α	IL8RA 2q34-q35	P55024	N.D.	None	None	N.A.	N.D.
CXCR2	IL8R _β , IL-8R ₂ , IL-8R _β	IL8RB 2q34-q35	P55025	P53543	GRX _α , NAP-2, RMO-78	SRB RS-145004-000 (35 nM)	Mice: Defective neutrophil trafficking and distribution and B cell distribution; resistance to adenovirus	N.D.
CXCR3	IP10/Mig R, GPR9	GPR9 Xq13	P45632	AF045146	IP-10, ITAC, Mig	None	N.A.	N.D.
CXCR4	HOMOXYN, LESTR, Isuin, HM80, LXA1, NYLA, D2850 IL	CXCR4 3p21-q21	P30991	P70656	SDP-1	AMD100	Mice: Defective development of heart, thymus, and bone marrow lymphopoiesis and bone marrow myelopoiesis Embryonic lethal (rescue by bone marrow transplantation)	X4 strains
CXCR5	BLR-1, MDR15	BLR1 11	P32202	Q04951	BCA-1	None	Mice: Defective B cell trafficking and distribution; thymic atrophy, Treg's patches and germinal centers	HIV 2
CC chemokine receptors								
CCL1	CKR1, CC CKR1, MIP-1α/RANTES, CMKR1	CCL1 3p21	P32246	P51675	HCC-1	Beda RS-162983-000 (6 nM)	Mice: Susceptibility to <i>A. fumigatus</i> , <i>S. aureus</i> , <i>Yersinia</i> abscessus, defective neutrophil chemotaxis, Th1/Th2 balance, hemopoiesis, and granuloma formation; susceptibility to neoplastic diseases; defective cardiac and chronic cardiac allograft rejection	N.D.
CCL2	CKR2, CC CKR2, CC CKR2, MCP-1, CMKR2	CCL2 3p21	P41597	P51683	MCP-1	Roche RS-511336-000 (33 nM)	Mice: Susceptibility to <i>Leishmania</i> , resistance to atypical mycobacteria, defective monocyte chemotaxis; Th1/Th2 balance, and granuloma formation	Some dual-tropic strains
CCL3	CKR3, CC CKR3, eotaxin receptor, CMKR3	CCL3 3p21	P51677	P51678	Eotaxin, eotaxin-2	Banya RS-163885-230 (40 nM)	N.A.	Some X4, R5, and dual-tropic strains
CCL4	CKR4, CC CKR4, R5-5, CMKR4, CCR4	CCL4 3p21	P51679	P51680	TARC, MDC	None	N.A.	N.D.
CCL5	CKR5, CC CKR5, ChemR13, CMKR5	CCL5 3p21	P51681	P51682	MIP-1β	Tkeda TAK-779 (1 nM)	N.A.	All R5 and dual-tropic strains
CCL6	GPR174, CKR14, STRL32, CCR6, CC CKR6, R5-1, GPR28, CMKR6	CCL6 6q27	P51684	AJ22714	LARC	None	N.A.	N.D.
CCL7	EBF-1, BLR-2, CMKR7	CCL7 17q12-q21.2	P32248	P4774	ELC, SLC	None	N.A.	N.D.
CCL8	TRE1, CKR11, GPR176, ChemR1, CMKR8	CCL8 3p21-q22	P51685	AF01277	L309	None	N.A.	Some X4, R5, and dual-tropic strains
CCL9	GPR 9-6	Unamed 3p21-q22	P51686	AJ13236	TECK	None	N.A.	N.D.
CCL10	GPR2	Unamed 17q21-q21.3	U13807		ESKine	None	N.A.	N.D.
CCL11	PTP1	GPR25 3p21	AF165507	N.A.	TBA	None	N.A.	N.D.
CXCR1 chemokine receptor	GPR5	GPR5 3p21	P46994	N.A.	Lymphotactin	None	N.A.	N.D.
CXCR3 chemokine receptor	GPR13, V28, CMKRBL1	CXCR3 3p21-q22	P46928	AF102269	Fractalkine	None	N.A.	Some X4, R5, and dual-tropic strains
Chemokine-binding proteins								
Duffy	DARC, glycosylated D	Fy	Q16570	AF016584	None	Yes	Human: Resistance to <i>P. vivax</i> malaria	N.D.
D6	CCR9, CCR10	CCL11-q22 CMAR9 3p21	Y12815	Y12815	None	N.A.	N.A.	N.D.

CX3C:	...CXXXC.....C.....C.....	n=1
CX:	...CX _C.....C.....C.....	n>15
CC:	...C _C.....C.....C.....	n>25
C:C.....C.....C.....	n=2

FIG. 1. Structural classification of the chemokine family by signature cysteines. The number of members in each subclass is listed at the right of each structure. Underlines indicate gaps in the alignment; X, an amino acid other than cysteine; and dots, other amino acids. Spacing between cysteines is similar in all four groups. The N and C termini can vary in length.

immune responses; a homeostatic/constitutive group, which is important in lymphocyte and dendritic cell trafficking in immune surveillance (Cyster, 1999a,b); and an overlap group. Genes encoding inflammatory chemokines are typically found in two major clusters on human chromosomes 4 (CXC) and 17 (CC), whereas genes for homeostatic chemokines are located alone or in small clusters on chromosomes 1, 2, 5, 7, 9, 10, and 16. Homeostatic receptors include CXCR4, CXCR5, CCR4, CCR7, and CCR9. Inflammatory receptors include CXCR1, CXCR2, CXCR3, CCR1, CCR2, CCR3, CCR5, and CCR6.

C. Chemokine Receptor Classification and Nomenclature

The classification of chemokine receptors is restricted to those defined at the molecular level. Native receptors are more difficult to study specifically because few selective agonists and antagonists are available and because multiple receptor subtypes with overlapping ligand specificities may be expressed in the same cell.

Although most chemokine receptors recognize more than one chemokine, they are almost always restricted to a single subclass (Table 4). Thus, the nomenclature system is rooted by the chemokine subclass specificity of the receptor. Human CC and CXC chemokine receptor names consist of the root CCR or CXCR, respectively, followed by a number. The lymphotactin and fractalkine receptors are named CXCR1 ["X" to distinguish it from complement receptor 1 (CR1)] and CX3CR1, respectively. The use of the letter "R" in receptor names is nonstandard for pharmacologists but is widely accepted practice for immunologists and was therefore authorized as an exception by NC-IUPHAR. Thus, these receptors are referred to as, for example, "CXCR1," and not "the CXCR1 receptor," which would be redundant. Splice variants, if pharmacologically distinct, are designated by a lowercase letter starting from the beginning of the alphabet subscripted in parentheses after the receptor name. Species orthologs are indicated by an appropriate species abbreviation followed by a space before the receptor name (Vanhoute et al., 1998). By consensus agreement of the conferees at the 1996 Gordon Conference on Chemotactic Cytokines, new names are assigned by a committee composed of Phil Murphy (pmm@nih.gov), Craig Gerard (gerard_c@

gonzo.tch.harvard.edu), and Tom Schall (tschall@chemocentryx.com).

Like the receptor names, systematic chemokine names, shown in Table 2 with their corresponding common names, are also built from cysteine subclass roots, followed by "L" for "ligand" and a number. In general, the numbers correspond to the same number used in the corresponding gene nomenclature, which takes the form "SCY" for "small cytokine," followed by "A," "B," "C," or "D" for "CC," "CXC," "C," or "CX3C" subclass, respectively, followed by the number.

Analysis of chemokine receptors presents problems not faced with other types of GPCRs. Most imposing is the large number of receptors and endogenous ligands and their overlapping specificities for each other and for leukocyte subtypes. In addition, both chemokines and their receptors may vary markedly in sequence among species, as much as 55% aa divergence in the case of certain chemokines. As a result, even though chemokine orthologs from different species usually cross-activate receptors, the receptors may have markedly different biology and pharmacology. Even the repertoire of chemokine and chemokine receptors may differ in different species. For example, mouse orthologs of IL-8 and CXCR1 have not been found, and there is persuasive evidence in the case of IL-8 that a mouse form does not exist (Modi and Yoshimura, 1999). Why these molecules are evolving so rapidly is unknown, but it is a property shared fairly selectively with the class of genes involved in immunity and inflammation (Murphy, 1993).

D. Chemokine Receptor Structure

The sequences of chemokine receptors have 25 to 80% aa identity (Fig. 2), indicating a common ancestor. However, many other G protein-coupled peptide receptors also have ~25% aa identity to chemokine receptors, illustrating that the structural boundary is not sharp. Although they lack a single structural signature, there are several features that together are found more frequently among chemokine receptors than other types of GPCRs. These include a length of 340 to 370 aa; an acidic N-terminal segment; the sequence DRYLAIVHA, or a variation of it, in the second intracellular loop; a short basic third intracellular loop; and a cysteine in each of the four extracellular domains. A tyrosine sulfation motif is commonly found in the N terminus of chemokine receptors and has been shown to be critical for HIV coreceptor activity for CCR5 (Farzan et al., 1999).

The three-dimensional structure of chemokine receptors is unknown, but a reasonable working model can be constructed for the transmembrane domains based on analogy with rhodopsin (Baldwin, 1993; Unger et al., 1997; Lomize et al., 1999). Models of the extracellular and intracellular domains are completely speculative, although in some cases domain-specific antibodies have verified the general location. Evidence has been reported that CCR2, CCR5, and CXCR4 form homodimers (Ben-




FIG. 2. Structural relationships among human chemokine receptors. The dendrogram was constructed by Marc Rothenberg using default parameters in the PILEUP algorithm of the University of Wisconsin Genetics Computer Group.

N-terminal segment is important in determining whether a given chemokine binds to receptor, and if so, whether it functions as an agonist or antagonist. Truncation may also cause a switch in receptor specificity as in the cases of NAP-2 and MCP-1.

E. Chemokine Receptor Specificity for Ligands and Leukocytes

Each chemokine receptor has a distinct chemokine and leukocyte specificity (Table 4), but the specificities can overlap considerably, because some chemokines can bind multiple receptor subtypes, and some receptors can bind multiple chemokines. Mutagenesis has indicated that the ligand binding site of chemokine receptors is highly complex, being composed of multiple noncontiguous domains and at least two distinct subsites: one for docking and the other for triggering (Ahuja et al., 1996; Monteclaro and Charo, 1996; Crump et al., 1997). At least for CCR5 and CXCR4, the first two TMDs and associated loops, but not the N-terminal segment, appear to be dispensable for normal receptor expression and function (Ling et al., 1999). Multiple low-affinity interactions together provide the high-affinity binding energy. A conserved HIV gp120 glycoprotein structure has been solved that is involved in chemokine receptor binding (Rizzuto et al., 1998).

Inflammatory chemokines (mainly those encoded by the chromosome 4 and 17 clusters of genes) have highly promiscuous relationships with receptors. There are fewer homeostatic chemokines, but those that map to the same chromosome tend to bind to the same receptor [e.g., MDC and TARC at CCR4; EBI ligand chemokine (ELC) and secondary lymphoid tissue chemokine (SLC) at CCR7]. Recently, the number of monogamous chemokine ligand-receptor relationships, which had previously been regarded as exceptional, has risen substantially (e.g., SDF-1 and CXCR4; TECK and CCR9; BLC and CXCR5; LARC and CCR6; lymphotactin and XCR1; fractalkine and CX3CR1).

Adding to the complexity of the system, distinct receptor subtypes specific for the same chemokine and the same function can be coexpressed on the same cell (Morohashi et al., 1995), distinct chemokines acting at separate receptors coexpressed on the same cell can induce the same cellular response (Zaitseva et al., 1997), and the same receptor can sort signals from different ligands to distinct signaling pathways (Zhang et al., 1999). Also, chemokine receptors are not limited to leukocytes but in specific cases may also be expressed on endothelial cells, neurons, epithelial cells, and microglial cells of the brain (Hadley et al., 1994; He et al., 1997; Horuk et al., 1997; Gupta et al., 1998b; Salcedo et al., 1999). There is intense interest in understanding the biological roles of these receptors in these ectopic sites.

With rare exceptions (Blanpain et al., 1999), functional human chemokines are agonists at leukocyte receptors. In contrast, naturally occurring chemokine antagonists have only been found in viruses (Table 5). For example, the viral chemokines MC148R from the poxvirus *Molluscum contagiosum* virus (Damon et al., 1998) and vMIP-II from human herpesvirus 8 (HHV8) (Kledal et al., 1997) are broad-spectrum chemokine receptor antagonists, suggesting roles in immune evasion and the importance of normal chemokine signaling for antiviral host defense. Various orthopoxviruses (e.g., myxoma, vaccinia) deploy an alternative strategy to block chemokines, through two structurally unique classes of secreted, broadly specific chemokine scavengers, one of which also binds interferon- γ (Graham et al., 1997; Smith et al., 1997; Alcami et al., 1998). Neither has structural homology to chemokines, chemokine receptors, or any other proteins currently in the public databases. They may be good leads for development of novel anti-inflammatory agents, particularly for topical or single-use administration (Table 5).

Recently, a growing number of structurally diverse, naturally occurring, nonchemokine ligands for chemokine receptors has been identified. These include HIV

TABLE 5

Viral chemokines and chemokine receptors
All molecules listed, except for the poxvirus chemokine binding proteins and HIV Tat, have conserved sequences with cellular chemokines or chemokine receptors (TTM). N.A., not available. Note that the following ORFs are syntenic: U12 of HHV6 and HHV7, M33 of MCMV, and R33 of rat CMV.

Virus Family	Virus	ORF	Common Name	Chemokine or Receptor Class	Function	References
γ -Herpesviridae	<i>H. simuli</i>	74	BCRP3	ELR+ CXCR	Calcium flux in vitro	Ahuja and Murphy, 1993
	HHV8 (KSHV)	74	ORF 74	CCXCR	Oncogenic	Arvanitakis et al., 1997
		K8	KSHV GPCR	(constitutively active)	Autogenic	Bae et al., 1996
		K4	vMIP-1	CC chemokine	Autogenic	Bohndorf et al., 1997
β -Herpesviridae			vMIP-1I	CC chemokine	Autogenic	Bohndorf et al., 1997
					THV suppressive	Kiehl et al., 1997
					TN α chemotactant	Soranzo et al., 1996b
					Antagonist at CXCR1, CXCR2, CXCR3, CXCR4, and CXCR5	Daugherty et al., 1999
β -Herpesviridae			vMIP-III	CC chemokine	N.A.	Virgin et al., 1997
			74	Potential CXCR	N.A.	Telford et al., 1995
				Potential CXCR	Chemotaxis	Camacho et al., 1999
				Potential CXCR	N.A.	Neale et al., 1998
			US28	CCXCR2	Chemokine sequestration	Gas and Murphy, 1994
					Smooth muscle cell migration	Bohndorf et al., 1999
					N.A.	Shedden et al., 1999
					Neutrophil calcium flux, chemotaxis, and degranulation; CXCR2 specific	Poulliat et al., 1999
					N.A.	Studerup et al., 1999
					Virulence factor. Blocks NK- and T cell response to MCMV in vivo; also poult-inflammatory early in infection	Plinning et al., 1999
Poxviridae			MCK-DMCK-2	CC chemokine	Calcium flux (mouse peritoneal macrophages; human CXCR3 transfectants)	Davis-Poyner et al., 1997
					Virulence factor	Reiser et al., 1998
					Virulence factor	Reiser et al., 1998
					Calcium flux in vitro	Reiser et al., 1998
					THP-1 cell chemotaxis	Zou et al., 1999
					Blocks neutrophil, monocyte, and T cell chemotaxis induced by multiple CC and CXCR chemokines	Nichols, 1996
					Antagonist at CXCR2 and CXCR3	Kratzwohl et al., 1997
					Induced by human hemopoietic progenitor cell proliferation	Donato et al., 1998
					Broad-spectrum CC chemokine scavenger	Daugherty et al., 1999
					Anti-inflammatory in context of vaccinia infection and allergic airway inflammation in guinea pig	Smith et al., 1997
Poxviridae				CC chemokine binding protein ($K_d = 0.1-15$ nM)	Broad-spectrum CC chemokine scavenger	Graham et al., 1997
					Anti-inflammatory in context of vaccinia infection and allergic airway inflammation in guinea pig	Smith et al., 1997
					Broad-spectrum CC chemokine and γ -IFN scavenger	Alcami et al., 1996
					Anti-inflammatory in context of myxoma infection	Lalani et al., 1997
Lentiviridae				N.A.	N.A.	Massing et al., 1993
					Monocyte chemotactant	Cao et al., 1995
					Antagonist at CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, CXCR7, CXCR8, CXCR9, CXCR10, CXCR11, CXCR12, CXCR13, CXCR14, CXCR15, CXCR16, CXCR17, CXCR18, CXCR19, CXCR20, CXCR21, CXCR22, CXCR23, CXCR24, CXCR25, CXCR26, CXCR27, CXCR28, CXCR29, CXCR30, CXCR31, CXCR32, CXCR33, CXCR34, CXCR35, CXCR36, CXCR37, CXCR38, CXCR39, CXCR40, CXCR41, CXCR42, CXCR43, CXCR44, CXCR45, CXCR46, CXCR47, CXCR48, CXCR49, CXCR50, CXCR51, CXCR52, CXCR53, CXCR54, CXCR55, CXCR56, CXCR57, CXCR58, CXCR59, CXCR60, CXCR61, CXCR62, CXCR63, CXCR64, CXCR65, CXCR66, CXCR67, CXCR68, CXCR69, CXCR70, CXCR71, CXCR72, CXCR73, CXCR74, CXCR75, CXCR76, CXCR77, CXCR78, CXCR79, CXCR80, CXCR81, CXCR82, CXCR83, CXCR84, CXCR85, CXCR86, CXCR87, CXCR88, CXCR89, CXCR90, CXCR91, CXCR92, CXCR93, CXCR94, CXCR95, CXCR96, CXCR97, CXCR98, CXCR99, CXCR100, CXCR101, CXCR102, CXCR103, CXCR104, CXCR105, CXCR106, CXCR107, CXCR108, CXCR109, CXCR110, CXCR111, CXCR112, CXCR113, CXCR114, CXCR115, CXCR116, CXCR117, CXCR118, CXCR119, CXCR120, CXCR121, CXCR122, CXCR123, CXCR124, CXCR125, CXCR126, CXCR127, CXCR128, CXCR129, CXCR130, CXCR131, CXCR132, CXCR133, CXCR134, CXCR135, CXCR136, CXCR137, CXCR138, CXCR139, CXCR140, CXCR141, CXCR142, CXCR143, CXCR144, CXCR145, CXCR146, CXCR147, CXCR148, CXCR149, CXCR150, CXCR151, CXCR152, CXCR153, CXCR154, CXCR155, CXCR156, CXCR157, CXCR158, CXCR159, CXCR160, CXCR161, CXCR162, CXCR163, CXCR164, CXCR165, CXCR166, CXCR167, CXCR168, CXCR169, CXCR170, CXCR171, CXCR172, CXCR173, CXCR174, CXCR175, CXCR176, CXCR177, CXCR178, CXCR179, CXCR180, CXCR181, CXCR182, CXCR183, CXCR184, CXCR185, CXCR186, CXCR187, CXCR188, CXCR189, CXCR190, CXCR191, CXCR192, CXCR193, CXCR194, CXCR195, CXCR196, CXCR197, CXCR198, CXCR199, CXCR200, CXCR201, CXCR202, CXCR203, CXCR204, CXCR205, CXCR206, CXCR207, CXCR208, CXCR209, CXCR210, CXCR211, CXCR212, CXCR213, CXCR214, CXCR215, CXCR216, CXCR217, CXCR218, CXCR219, CXCR220, CXCR221, CXCR222, CXCR223, CXCR224, CXCR225, CXCR226, CXCR227, CXCR228, CXCR229, CXCR230, CXCR231, CXCR232, CXCR233, CXCR234, CXCR235, CXCR236, CXCR237, CXCR238, CXCR239, CXCR240, CXCR241, CXCR242, CXCR243, CXCR244, CXCR245, CXCR246, CXCR247, CXCR248, CXCR249, CXCR250, CXCR251, CXCR252, CXCR253, CXCR254, CXCR255, CXCR256, CXCR257, CXCR258, CXCR259, CXCR260, CXCR261, CXCR262, CXCR263, CXCR264, CXCR265, CXCR266, CXCR267, CXCR268, CXCR269, CXCR270, CXCR271, CXCR272, CXCR273, CXCR274, CXCR275, CXCR276, CXCR277, CXCR278, CXCR279, CXCR280, CXCR281, CXCR282, CXCR283, CXCR284, CXCR285, CXCR286, CXCR287, CXCR288, CXCR289, CXCR290, CXCR291, CXCR292, CXCR293, CXCR294, CXCR295, CXCR296, CXCR297, CXCR298, CXCR299, CXCR300, CXCR301, CXCR302, CXCR303, CXCR304, CXCR305, CXCR306, CXCR307, CXCR308, CXCR309, CXCR310, CXCR311, CXCR312, CXCR313, CXCR314, CXCR315, CXCR316, CXCR317, CXCR318, CXCR319, CXCR320, CXCR321, CXCR322, CXCR323, CXCR324, CXCR325, CXCR326, CXCR327, CXCR328, CXCR329, CXCR330, CXCR331, CXCR332, CXCR333, CXCR334, CXCR335, CXCR336, CXCR337, CXCR338, CXCR339, CXCR340, CXCR341, CXCR342, CXCR343, CXCR344, CXCR345, CXCR346, CXCR347, CXCR348, CXCR349, CXCR350, CXCR351, CXCR352, CXCR353, CXCR354, CXCR355, CXCR356, CXCR357, CXCR358, CXCR359, CXCR360, CXCR361, CXCR362, CXCR363, CXCR364, CXCR365, CXCR366, CXCR367, CXCR368, CXCR369, CXCR370, CXCR371, CXCR372, CXCR373, CXCR374, CXCR375, CXCR376, CXCR377, CXCR378, CXCR379, CXCR380, CXCR381, CXCR382, CXCR383, CXCR384, CXCR385, CXCR386, CXCR387, CXCR388, CXCR389, CXCR390, CXCR391, CXCR392, CXCR393, CXCR394, CXCR395, CXCR396, CXCR397, CXCR398, CXCR399, CXCR400, CXCR401, CXCR402, CXCR403, CXCR404, CXCR405, CXCR406, CXCR407, CXCR408, CXCR409, CXCR410, CXCR411, CXCR412, CXCR413, CXCR414, CXCR415, CXCR416, CXCR417, CXCR418, CXCR419, CXCR420, CXCR421, CXCR422, CXCR423, CXCR424, CXCR425, CXCR426, CXCR427, CXCR428, CXCR429, CXCR430, CXCR431, CXCR432, CXCR433, CXCR434, CXCR435, CXCR436, CXCR437, CXCR438, CXCR439, CXCR440, CXCR441, CXCR442, CXCR443, CXCR444, CXCR445, CXCR446, CXCR447, CXCR448, CXCR449, CXCR450, CXCR451, CXCR452, CXCR453, CXCR454, CXCR455, CXCR456, CXCR457, CXCR458, CXCR459, CXCR460, CXCR461, CXCR462, CXCR463, CXCR464, CXCR465, CXCR466, CXCR467, CXCR468, CXCR469, CXCR470, CXCR471, CXCR472, CXCR473, CXCR474, CXCR475, CXCR476, CXCR477, CXCR478, CXCR479, CXCR480, CXCR481, CXCR482, CXCR483, CXCR484, CXCR485, CXCR486, CXCR487, CXCR488, CXCR489, CXCR490, CXCR491, CXCR492, CXCR493, CXCR494, CXCR495, CXCR496, CXCR497, CXCR498, CXCR499, CXCR500, CXCR501, CXCR502, CXCR503, CXCR504, CXCR505, CXCR506, CXCR507, CXCR508, CXCR509, CXCR510, CXCR511, CXCR512, CXCR513, CXCR514, CXCR515, CXCR516, CXCR517, CXCR518, CXCR519, CXCR520, CXCR521, CXCR522, CXCR523, CXCR524, CXCR525, CXCR526, CXCR527, CXCR528, CXCR529, CXCR530, CXCR531, CXCR532, CXCR533, CXCR534, CXCR535, CXCR536, CXCR537, CXCR538, CXCR539, CXCR540, CXCR541, CXCR542, CXCR543, CXCR544, CXCR545, CXCR546, CXCR547, CXCR548, CXCR549, CXCR550, CXCR551, CXCR552, CXCR553, CXCR554, CXCR555, CXCR556, CXCR557, CXCR558, CXCR559, CXCR560, CXCR561, CXCR562, CXCR563, CXCR564, CXCR565, CXCR566, CXCR567, CXCR568, CXCR569, CXCR570, CXCR571, CXCR572, CXCR573, CXCR574, CXCR575, CXCR576, CXCR577, CXCR578, CXCR579, CXCR580, CXCR581, CXCR582, CXCR583, CXCR584, CXCR585, CXCR586, CXCR587, CXCR588, CXCR589, CXCR590, CXCR591, CXCR592, CXCR593, CXCR594, CXCR595, CXCR596, CXCR597, CXCR598, CXCR599, CXCR600, CXCR601, CXCR602, CXCR603, CXCR604, CXCR605, CXCR606, CXCR607, CXCR608, CXCR609, CXCR610, CXCR611, CXCR612, CXCR613, CXCR614, CXCR615, CXCR616, CXCR617, CXCR618, CXCR619, CXCR620, CXCR621, CXCR622, CXCR623, CXCR624, CXCR625, CXCR626, CXCR627, CXCR628, CXCR629, CXCR630, CXCR631, CXCR632, CXCR633, CXCR634, CXCR635, CXCR636, CXCR637, CXCR638, CXCR639, CXCR640, CXCR641, CXCR642, CXCR643, CXCR644, CXCR645, CXCR646, CXCR647, CXCR648, CXCR649, CXCR650, CXCR651, CXCR652, CXCR653, CXCR654, CXCR655, CXCR656, CXCR657, CXCR658, CXCR659, CXCR660, CXCR661, CXCR662, CXCR663, CXCR664, CXCR665, CXCR666, CXCR667, CXCR668, CXCR669, CXCR670, CXCR671, CXCR672, CXCR673, CXCR674, CXCR675, CXCR676, CXCR677, CXCR678, CXCR679, CXCR680, CXCR681, CXCR682, CXCR683, CXCR684, CXCR685, CXCR686, CXCR687, CXCR688, CXCR689, CXCR690, CXCR691, CXCR692, CXCR693, CXCR694, CXCR695, CXCR696, CXCR697, CXCR698, CXCR699, CXCR700, CXCR701, CXCR702, CXCR703, CXCR704, CXCR705, CXCR706, CXCR707, CXCR708, CXCR709, CXCR710, CXCR711, CXCR712, CXCR713, CXCR714, CXCR715, CXCR716, CXCR717, CXCR718, CXCR719, CXCR720, CXCR721, CXCR722, CXCR723, CXCR724, CXCR725, CXCR726, CXCR727, CXCR728, CXCR729, CXCR730, CXCR731, CXCR732, CXCR733, CXCR734, CXCR735, CXCR736, CXCR737, CXCR738, CXCR739, CXCR740, CXCR741, CXCR742, CXCR743, CXCR744, CXCR745, CXCR746, CXCR747, CXCR748, CXCR749, CXCR750, CXCR751, CXCR752, CXCR753, CXCR754, CXCR755, CXCR756, CXCR757, CXCR758, CXCR759, CXCR760, CXCR761, CXCR762, CXCR763, CXCR764, CXCR765, CXCR766, CXCR767, CXCR768, CXCR769, CXCR770, CXCR771, CXCR772, CXCR773, CXCR774, CXCR775, CXCR776, CXCR777, CXCR778, CXCR779, CXCR780, CXCR781, CXCR782, CXCR783, CXCR784, CXCR785, CXCR786, CXCR787, CXCR788, CXCR789, CXCR790, CXCR791, CXCR792, CXCR793, CXCR794, CXCR795, CXCR796, CXCR797, CXCR798, CXCR799, CXCR800, CXCR801, CXCR802, CXCR803, CXCR804, CXCR805, CXCR806, CXCR807, CXCR808, CXCR809, CXCR810, CXCR811, CXCR812, CXCR813, CXCR814, CXCR815, CXCR816, CXCR817, CXCR818, CXCR819, CXCR820, CXCR821, CXCR822, CXCR823, CXCR824, CXCR825, CXCR826, CXCR827, CXCR828, CXCR829, CXCR830, CXCR831, CXCR832, CXCR833, CXCR834, CXCR835, CXCR836, CXCR837, CXCR838, CXCR839, CXCR840, CXCR841, CXCR842, CXCR843, CXCR844, CXCR845, CXCR846, CXCR847, CXCR848, CXCR849, CXCR850, CXCR851, CXCR852, CXCR853, CXCR854, CXCR855, CXCR856, CXCR857, CXCR858, CXCR859, CXCR860, CXCR861, CXCR862, CXCR863, CXCR864, CXCR865, CXCR866, CXCR867, CXCR868, CXCR869, CXCR870, CXCR871, CXCR872, CXCR873, CXCR874, CXCR875, CXCR876, CXCR877, CXCR878, CXCR879, CXCR880, CXCR881, CXCR882, CXCR883, CXCR884, CXCR885, CXCR886, CXCR887, CXCR888, CXCR889, CXCR890, CXCR891, CXCR892, CXCR893, CXCR894, CXCR895, CXCR896, CXCR897, CXCR898, CXCR899, CXCR900, CXCR901, CXCR902, CXCR903, CXCR904, CXCR905, CXCR906, CXCR907, CXCR908, CXCR909, CXCR910, CXCR911, CXCR912, CXCR913, CXCR914, CXCR915, CXCR916, CXCR917, CXCR918, CXCR919, CXCR920, CXCR921, CXCR922, CXCR923, CXCR924, CXCR925, CXCR926, CXCR927, CXCR928, CXCR929, CXCR930, CXCR931, CXCR932, CXCR933, CXCR934, CXCR935, CXCR936, CXCR937, CXCR938, CXCR939, CXCR940, CXCR941, CXCR942, CXCR943, CXCR944, CXCR945, CXCR946, CXCR947, CXCR948, CXCR949, CXCR950, CXCR951, CXCR952, CXCR953, CXCR954, CXCR955, CXCR956, CXCR957, CXCR958, CXCR959, CXCR960, CXCR961, CXCR962, CXCR963, CXCR964, CXCR965, CXCR966, CXCR967, CXCR968, CXCR969, CXCR970, CXCR971, CXCR972, CXCR973, CXCR974, CXCR975, CXCR976, CXCR977, CXCR978, CXCR979, CXCR980, CXCR981, CXCR982, CXCR983, CXCR984, CXCR985, CXCR986, CXCR987, CXCR988, CXCR989, CXCR990, CXCR991, CXCR992, CXCR993, CXCR994, CXCR995, CXCR996, CXCR997, CXCR998, CXCR999, CXCR1000, CXCR1001, CXCR1002, CXCR1003, CXCR1004, CXCR1005, CXCR1006, CXCR1007, CXCR1008, CXCR1009, CXCR1010, CXCR1011, CXCR1012, CXCR1013, CXCR1014, CXCR1015, CXCR1016, CXCR1017, CXCR1018, CXCR1019, CXCR1020, CXCR1021, CXCR1022, CXCR1023, CXCR1024, CXCR1025, CXCR1026, CXCR1027, CXCR1028, CXCR1029, CXCR1030, CXCR1031, CXCR1032, CXCR1033, CXCR1034, CXCR1035, CXCR1036, CXCR1037, CXCR1038, CXCR1039, CXCR1040, CXCR1041, CXCR1042, CXCR1043, CXCR1044, CXCR1045, CXCR1046, CXCR1047, CXCR1048, CXCR1049, CXCR1050, CXCR1051, CXCR1052, CXCR1053, CXCR1054, CXCR1055, CXCR1056, CXCR1057, CXCR1058, CXCR1059, CXCR1060, CXCR1061, CXCR1062, CXCR1063, CXCR1064, CXCR1065, CXCR1066, CXCR1067, CXCR1068, CXCR1069, CXCR1070, CXCR1071, CXCR1072, CXCR1073, CXCR1074, CXCR1075, CXCR1076, CXCR1077, CXCR1078, CXCR1079, CXCR1080, CXCR1081, CXCR1082, CXCR1083, CXCR1084, CXCR1085, CXCR1086, CXCR1087, CXCR1088, CXCR1089, CXCR1090, CXCR1091, CXCR1092, CXCR1093, CXCR1094, CXCR1095, CXCR1096, CXCR1097, CXCR1098, CXCR1099, CXCR1100, CXCR1101, CXCR1102, CXCR1103, CXCR1104, CXCR1105, CXCR1106, CXCR1107, CXCR1108, CXCR1109, CXCR1110, CXCR1111, CXCR1112, CXCR1113, CXCR1114, CXCR1115, CXCR1116, CXCR1117, CXCR1118, CXCR1119, CXCR1120, CXCR1121, CXCR1122, CXCR1123, CXCR1124, CXCR1125, CXCR1126, CXCR1127, CXCR1128, CXCR1129, CXCR1130, CXCR1131, CXCR1132, CXCR1133, CXCR1134, CXCR1135, CXCR1136, CXCR1137, CXCR1138, CXCR1139, CXCR1140, CXCR1141, CXCR1142, CXCR1143, CXCR1144, CXCR1145, CXCR1146, CXCR1147, CXCR1148, CXCR1149, CXCR1150, CXCR1151, CXCR1152, CXCR1153, CXCR1154, CXCR1155, CXCR1156, CXCR1157, CXCR1158, CXCR1159, CXCR1160, CXCR1161, CXCR1162, CXCR1163, CXCR1164, CXCR1165, CXCR1166, CXCR1167, CXCR1168, CXCR1169, CXCR1170, CXCR1171, CXCR1172, CXCR1173, CXCR1174, CXCR1175, CXCR1176, CXCR1177, CXCR1178, CXCR1179, CXCR1180, CXCR1181, CXCR1182, CXCR1183, CXCR1184, CXCR1185, CXCR1186, CXCR1187, CXCR1188, CXCR1189, CXCR1190, CXCR1191, CXCR1192, CXCR1193, CXCR1194, CXCR1195, CXCR1196, CXCR1197, CXCR1198, CXCR1199, CXCR1200, CXCR1201, CXCR1202, CXCR1203, CXCR1204, CXCR1205, CXCR1206, CXCR1207, CXCR1208, CXCR1209, CXCR1210, CXCR1211, CXCR1212, CXCR1213, CXCR1214, CXCR1215, CXCR1216, CXCR1217, CXCR1218, CXCR1219, CXCR1220, CXCR1221, CXCR1222, CXCR1223, CXCR1224, CXCR1225, CXCR1226, CXCR1227, CXCR1228, CXCR1229, CXCR1230, CXCR1231, CXCR1232, CXCR1233, CXCR1234, CXCR1235, CXCR1236, CXCR1237, CXCR1238, CXCR1239, CXCR1240, CXCR1241, CXCR1242, CXCR1243, CXCR1244, CXCR1245, CXCR1246, CXCR1247, CXCR1248, CXCR1249, CXCR1250, CXCR1251, CXCR1252, CXCR1253, CXCR1254, CXCR1255, CXCR1256, CXCR1257, CXCR1258, CXCR1259, CXCR1260, CXCR1261, CXCR1262, CXCR1263, CXCR1264, CXCR1265, CXCR1266, CXCR1267, CXCR1268, CXCR1269, CXCR1270, CXCR1271, CXCR1272, CXCR1273, CXCR1274, CXCR1275, CXCR1276, CXCR1277, CXCR1278, CXCR1279, CXCR1280, CXCR1281, CXCR1282, CXCR1283, CXCR1284, CXCR1285, CXCR1286, CXCR1287, CXCR1288, CXCR1289, CXCR1290, CXCR1291, CXCR1292, CXCR1293, CXCR1294, CXCR1295, CXCR1296, CXCR1297, CXCR1298, CXCR1299, CXCR1300, CXCR1301, CXCR1302, CXCR1303, CXCR1304, CXCR1305, CXCR1306, CXCR1307, CXCR1308, CXCR1309, CXCR1310, CXCR1311, CXCR1312, CXCR1313, CXCR1314, CXCR1315, CXCR1316, CXCR1317, CXCR1318, CXCR1319, CXCR1320, CXCR1321, CXCR1322, CXCR1323, CXCR1324, CXCR1325, CXCR1326, CXCR1327, CXCR1328, CXCR1329, CXCR1330, CXCR1331, CXCR1332, CXCR1333, CXCR1334, CXCR1335, CXCR1336, CXCR1337, CXCR1338, CXCR1339, CXCR1340, CXCR1341, CXCR1342, CXCR1343, CXCR1344, CXCR1345, CXCR1346, CXCR1347, CXCR1348, CXCR1349, CXCR1350, CXCR1351, CXCR1352, CXCR1353, CXCR1354, CXCR1355, CXCR1356, CXCR1357, CXCR1358, CXCR1359, CXCR1360, CXCR1361, CXCR1362, CXCR1363, CXCR1364, CXCR1365, CXCR1366, CXCR1367, CXCR1368, CXCR1369, CXCR1370, CXCR1371, CXCR1372, CXCR1373, CXCR1374, CXCR1375, CXCR1376, CXCR1377, CXCR1378, CXCR1379, CXCR1380, CXCR1381, CXCR1382, CXCR1383, CXCR1384, CXCR1385, CXCR1386, CXCR1387, CXCR1388, CXCR1389, CXCR1390, CXCR1391, CXCR1392, CXCR1393, CXCR1394, CXCR1395, CXCR1396, CXCR1397, CXCR1398, CXCR1399, CXCR1400, CXCR1401, CXCR1402, CXCR1403, CXCR1404, CXCR1405, CXCR1406, CXCR1407, CXCR1408, CXCR1409, CXCR1410, CXCR1411, CXCR1412, CXCR1413, CXCR1414, CXCR1415, CXCR1416, CXCR1417, CXCR1418, CXCR1419, CXCR1420, CXCR1421, CXCR1422, CXCR1423, CXCR1424, CXCR1425, CXCR1426, CXCR1427, CXCR1428, CXCR1429, CXCR1430, CXCR1431, CXCR1432, CXCR1433, CXCR1434, CXCR1435, CXCR1436, CXCR1437, CXCR1438, CXCR1439, CXCR1440, CXCR1441, CXCR1442, CXCR1443, CXCR1444, CXCR1445, CXCR1446, CXCR1447, CXCR1448, CXCR1449, CXCR1450, CXCR1451, CXCR1452, CXCR1453, CXCR1454, CXCR1455, CXCR1456, CXCR1457, CXCR1458, CXCR1459, CXCR1460, CXCR1461, CXCR1462, CXCR1463, CXCR1464, CXCR1465, CXCR1466, CXCR1467, CXCR1468, CXCR1469, CXCR1470, CXCR1471, CXCR1472, CXCR1473, CXCR1474, CXCR1475, CXCR1476, CXCR1477, CXCR1478, CXCR1479, CXCR1480, CXCR1481, CXCR1482, CXCR1483, CXCR1484, CXCR1485, CXCR1486, CXCR1487, CXCR1488, CXCR1489, CXCR1490, CXCR1491, CXCR1492, CXCR1493, CXCR1494, CXCR1495, CXCR1496, CXCR1497, CXCR1498, CXCR1499, CXCR1500, CXCR1501, CXCR1502, CXCR1503, CXCR1504, CXCR1505, CXCR1506, CXCR1507, CXCR1508, CXCR1509, CXCR1510, CXCR1511, CXCR1512, CXCR1513, CXCR1514, CXCR1515, CXCR1516, CXCR1517, CXCR1518, CXCR1519, CXCR1520, CXCR1521, CXCR1522, CXCR1523, CXCR1524, CXCR1525, CXCR1526, CXCR1527, CXCR1528, CXCR1529, CXCR1530, CXCR1531, CXCR1532, CXCR1533, CXCR1534, CXCR1535, CXCR1536, CXCR1537, CXCR1538, CXCR1539, CXCR1540, CXCR1541, CXCR1542, CXCR1543, CXCR1544, CXCR1545, CXCR1546, CXCR1547, CXCR1548, CXCR1549, CXCR1550, CXCR1551, CXCR1552, CXCR1553, CXCR1554, CXCR1555, CXCR1556, CXCR1557, CXCR1558, CXCR1559, CXCR1560, CXCR1561, CXCR1562, CXCR1563, CXCR1564, CXCR1565, CXCR1566, CXCR1567, CXCR1568, CXCR1569, CXCR1570	

tat at CCR2 (Albini et al., 1998) and CXCR4 (Xiao et al., submitted), HIV gp120 at various HIV coreceptors (Berger et al., 1999), a secreted domain of tyrosyl tRNA synthetase at CXCR1 (Wakasugi et al., 1999), and the human β -defensin HBD2 at CCR6 (Yang et al., 1999).

A major new concept to emerge recently from studies of the leukocyte selectivities of chemokines is that interaction between antigen-loaded dendritic cells and antigen-specific T cells to achieve proper cell positioning in the periphery or in secondary lymphoid tissue for an adaptive immune response is not random but instead results in part from dynamic and coordinated changes in chemokine receptor expression. Moreover, the nature and strength of the immune response may be governed in part by specific chemokine receptor expression patterns. Thus, T lymphocytes and dendritic cells undergo highly dynamic regulation of chemokine receptors depending on whether the T cell is naive or memory, Th1 or Th2, and resting or activated, and whether the dendritic cell is immature or mature (Sallusto et al., 1999a; Sozzani et al., 1999). For example, when naive T lymphocytes are activated and differentiate into memory/effector cells, they down-regulate receptors for constitutive chemokines such as CXCR4 and CCR7 and acquire receptors for inflammatory chemokines such as CCR3, CCR5, and CXCR3. Also, dendritic cell maturation after antigen loading is accompanied by a transition from expression of inflammatory to homeostatic chemokine receptors. Distinct selectivities for Th1 and Th2 polarized T lymphocytes have been reported for CC chemokine receptors, and actual chemokine receptor markers of these cell types have been claimed and debated (Sallusto et al., 1998; Annunziato et al., 1999). Moreover, homing of memory T cells to specific anatomic sites has been strongly correlated with specific chemokine receptor expression patterns (Campbell et al., 1999).

With this as a general introduction, the next sections are discussions of the molecular pharmacology and biology of individual chemokine receptor subtypes. Note that the voluminous literature correlating the presence of specific chemokines in disease has been extensively reviewed (Baggiolini et al., 1997) and is not repeated here. Instead, we emphasize direct tests of function of specific receptors in disease.

III. CXC Chemokine Receptor Subtypes

A. CXCR1 and CXCR2

CXCR1 and CXCR2 were the first chemokine receptor subtypes to be defined. They are the only known mammalian receptors for ELR+ CXC chemokines, including IL-8, which binds to both receptors with similar high affinity; they do not bind other types of chemokines. They are also the major chemokine receptors expressed on neutrophils and are prototypic receptors for inflammatory/inducible chemokines. They appear to operate mainly in acute inflammation and innate immunity al-

though a role in macrophage accumulation in atherosclerotic plaque has also been demonstrated for CXCR2 (Boisvert et al., 1998). They are considered together because of these shared properties.

CXCR1 cDNA was first cloned from rabbit neutrophils by homology hybridization using a probe based on conserved sequences in TMD 2 of known neuropeptide-specific GPCRs (Thomas et al., 1990). When expressed in frog oocytes, it appeared to be specific for formyl-methionyl-leucyl-phenylalanine, but this could not be reproduced in mammalian cells where IL-8 was a functional ligand (Thomas et al., 1991). Consistent with this, human CXCR1 cDNA was isolated independently from a neutrophil library by expression cloning using an 125 I-IL-8 binding assay in COS-7 cells (Holmes et al., 1991). CXCR2 cDNA was first cloned by homology hybridization from a dibutyryl cAMP-induced HL-60 cell library using an oligonucleotide probe corresponding to TMD2 of rabbit CXCR1 (Murphy and Tiffany, 1991); later, cDNAs were also isolated from a neutrophil library (Lee et al., 1992). The genes, designated *i18ra* and *i18rb*, are located 20 kb apart on human chromosome 2q35, and there is a linked pseudogene of CXCR2 named *i18rp* (Ahuja et al., 1992; White et al., 1994). The open reading frames (ORFs), which each occupy a single exon, are 350 codons for CXCR1 and either 355 or 360 codons for CXCR2 (both of two in-frame ATG codons are flanked by favorable Kozak sequences). CXCR1 and CXCR2 are 78% identical in aa sequence.

In addition to neutrophils and monocyte/macrophages, CXCR1 and CXCR2 have been detected on cytokine-activated eosinophils, basophils, T lymphocytes, mast cells, and dendritic cells, but important functional roles in vivo have not been clearly demonstrated (Chuntharapai et al., 1994; Hammond et al., 1995; Morohashi et al., 1995; Xu et al., 1995; Heath et al., 1997; Sozzani et al., 1997; Nilsson et al., 1999; Ochensberger et al., 1999; Petering et al., 1999). CXCR2, but not CXCR1, has been identified on brain Purkinje cells by mAb and radioligand binding, but function remains undefined there as well (Horuk et al., 1997).

In calcium flux and chemotaxis assays, CXCR2 is relatively nonselective for IL-8 versus all other ELR+ CXC chemokines studied (<10-fold range in EC_{50}), whereas CXCR1 is highly selective for IL-8 (>50-fold difference in EC_{50}) (Lee et al., 1992; Loetscher et al., 1994; Ahuja and Murphy, 1996). GCP-2 is an equipotent agonist at both CXCR1 and CXCR2 (Wuyts et al., 1997). Thus, GRO α , NAP-2, and ENA-78 are selective ligands for CXCR2. Recently, a selective nonchemokine endogenous ligand was identified for CXCR1: the N-terminal cytokine module of human tyrosyl tRNA synthetase, which contains an ELR motif and functions as a neutrophil chemoattractant in vitro. Its biological function is not established but could involve inflammatory signaling by apoptotic cells (Wakasugi and Schimmel, 1999). Consistent with coexpression of CXCR1 and CXCR2 on neutro-

phils, IL-8 effectively blocks binding of other ELR+ CXC radioligands to human neutrophils and interferes with signaling (calcium flux), but conversely, other ELR+ CXC chemokines can only partially block IL-8 binding to neutrophils and subsequent calcium flux (Moser et al., 1991; Ahuja and Murphy, 1996). The receptors appear to function independently.

Antagonists at CXCR2 include N-terminal truncations of IL-8 and GRO α (Hesselgesser et al., 1995), selective neutralizing monoclonal and polyclonal antibodies (Hammond et al., 1995; Green et al., 1996; Jones et al., 1996), a small peptide of undefined selectivity (Hayashi et al., 1995), and SB 225002 [N-(2-hydroxy-4-nitrophenyl)-N'-(2-bromophenyl)urea], a selective small molecule, nonpeptide inhibitor of CXCR2 (White et al., 1998) (Fig. 3). The latter is a potent antagonist of 125 I-IL-8 binding with an IC_{50} value of 22 nM and has >150-fold selectivity over CXCR1. In vitro, SB 225002 potently inhibits human and rabbit neutrophil chemotaxis induced by both IL-8 and GRO α , and in vivo it selectively blocks IL-8-induced neutrophil margination in rabbits.

In vivo roles of IL-8 and related ligands have been extensively studied, but specific receptor roles are less well defined. The mouse chemokines MIP-2 and KC are human GRO homologs specific for mouse CXCR2 (Bozic et al., 1994; Lee et al., 1995). CXCR2 knockout mice fail to mobilize neutrophils to chemically irritated peritoneum in vivo, and $-/-$ neutrophils do not migrate in vitro in response to KC or MIP-2, indicating that CXCR2

is the dominant neutrophil receptor for these chemokines (Cacalano et al., 1994).

Unexpectedly, CXCR2 $-/-$ mice have massive expansion of neutrophils and B cells throughout the hematopoietic system when derived in specific pathogen-free conditions but not in germ-free conditions (Moore et al., 1995). The explanation may reside in part in the fact that CXCR2 is a negative regulator of hematopoiesis (Broxmeyer et al., 1996). Alternatively, Cacalano et al. (1994) speculated that the inability to properly survey tissues and eliminate external pathogens in the knockouts may result in the release of cytokines that stimulate neutrophil and B cell production. However, the animals have not been reported to have increased susceptibility to infectious disease from either environmental or challenge pathogens.

The defect in neutrophil-mediated inflammation in these mice is consistent with the effects of CXCR2 ligand neutralization in mouse (KC, MIP-2) and rabbit (IL-8) in diverse models of acute inflammation (skin, airway, pleura, glomeruli) (Sekido et al., 1993; e.g., Broaddus et al., 1994). These results suggest indications for IL-8 receptor antagonists in diseases such as psoriasis, coronary artery reperfusion injury, and acute glomerulonephritis. Still, it is important to point out that rodents are poor models of the human IL-8 signaling system: they lack IL-8, a mouse counterpart of CXCR1 has not been identified, and rat CXCR1 is expressed in macrophages not neutrophils (Dunstan et al., 1996). Nevertheless,




FIG. 3. Structures of nonpeptide small molecule antagonists of specific chemokine receptors. Note that a basic nitrogen is a common feature. (Figure courtesy of Kurt Jarnagin.)

IL-8 receptor function in the monocyte/macrophage lineage may be more important than was initially appreciated. In particular, IL-8 can trigger firm adhesion of human monocytes to vascular endothelium under flow conditions (Gerszten et al., 1999), and CXCR2 is critical for macrophage accumulation in atherosclerotic lesions of LDL receptor-deficient mice (Boisvert et al., 1998).

CXCR1 and CXCR2 have been reported to carry out different functional roles in human neutrophils *in vitro*. CXCR1 appears to be dominant for chemotaxis, superoxide production, and phospholipase D activation in response to IL-8 (Hammond et al., 1995; Jones et al., 1996), as well as for chemotaxis to NAP-2 at high concentrations ($>1 \mu\text{M}$) (Ludwig et al., 1997), whereas CXCR2 appears to mediate neutrophil chemotaxis to NAP-2 (and GRO α) at low concentrations. Calcium flux and degranulation are mediated through both receptors. However, cell migration may be more important than cell activation for IL-8 receptor function *in vivo*, as suggested by the accumulation of unactivated neutrophils and the lack of inflammatory pathology at sites of KC transgene expression in mice; this may be a general property of chemokines (Lira et al., 1994).

Despite abundant evidence that IL-8 is important in acute inflammation, proof of concept is still lacking for differential roles of CXCR1 versus CXCR2 *in vivo* and in human disease. As suggested earlier, major obstacles include the inadequacy of small animal models and the lack of adequate selective small molecule antagonists. Other major unanswered questions about these receptors include their structure, the relative roles of CXCR1 and CXCR2 in ELR+ CXC chemokine-induced angiogenesis and modulation of myelopoiesis (Brommeyer et al., 1997), and the putative function of CXCR2 in brain (Horuk et al., 1997).

Two functional viral homologs of CXCR2 have been identified, ECRF3 of Herpesvirus saimiri (Ahuja et al., 1993) and KSHV GPCR of KSHV (HHV8) (Arvanitakis et al., 1997), which are quite different from CXCR2 and are reviewed in a later section.

B. CXCR3

CXCR3 is the first chemokine receptor identified that is highly induced by T cell activation. The ORF was first identified in incomplete form in 1995 on a genomic clone isolated by polymerase chain reaction-based homology hybridization. The gene was named *GPR9* and was originally mapped incorrectly to human chromosome 8p11.2-12 (Marchese et al., 1995) and later mapped correctly to Xq13 (Loetscher et al., 1998a). A full-length cDNA was independently isolated from an IL-2-activated T cell library (Loetscher et al., 1996). The ORF is interrupted by one intron in the region encoding the N-terminal segment and predicts a polypeptide 368 aa in length. The deduced protein sequence of human CXCR3 is ~30% identical with CXCR1 and CXCR2.

CXCR3 binds three highly potent, inflammatory/inducible, ELR-negative CXC chemokine agonists, I-TAC, Mig, and IP-10 (Loetscher et al., 1998a; Cole et al., 1998; Weng et al., 1998), all of which chemoattract and induce calcium flux in activated T cells, tumor-infiltrating lymphocytes, and CXCR3-transfected cells. The rank order of binding affinity is I-TAC $>$ Mig \sim IP-10. Curiously, the human CC chemokines eotaxin and MCP-4 also bind to CXCR3-transfected cells but with much lower affinity ($K_i \sim 60 \text{ nM}$) and without activating the receptor (Weng et al., 1998). Also, the mouse CC chemokine SLC/6CKine has been reported to induce calcium flux through mouse CXCR3 (Soto et al., 1998), but this was not observed with human 6CKine with either human or mouse CXCR3 (87% aa identity) (Jenah et al., 1999). A CXCR3-specific mAb named 1C6 has been reported that blocks human IP-10, but not human Mig, binding to CXCR3 (Qin et al., 1998).

CXCR3 is expressed on a portion of circulating blood T cells, B cells, and natural killer (NK) cells (Qin et al., 1998). Although freshly isolated T cells respond to Mig, curiously they are relatively less responsive to IP-10. Expression and responsiveness are both markedly increased by T cell activation (Rabin et al., 1999), classifying CXCR3 as an inflammatory/inducible type of chemokine receptor. CXCR3 has been detected preferentially on Th1 T cell lines and clones *in vitro* but could not discriminate between Th1- (Crohn's disease) and Th2- (systemic sclerosis) dominant responses *in vivo* and therefore may not be a practical marker of Th1 cells, as had been suggested (Bonecchi et al., 1998; Sallusto et al., 1998, 1999b; Annunziati et al., 1999). Blood T cells expressing CXCR3 are mostly CD45RO+ memory cells and express high levels of $\beta 1$ -integrins. Virtually all T cells in rheumatoid arthritis synovial fluid and in various inflamed tissues, such as in ulcerative colitis, chronic vaginitis, and sarcoidosis, express CXCR3, particularly in perivascular regions, whereas fewer T cells within normal lymph nodes are positive (Agostini et al., 1998; Qin et al., 1998). CXCR3 is also consistently detected in functional form on transformed B cells from CLL patients (Trentin et al., 1999).

The biological role of CXCR3 is not yet known, and it has not been established as a disease target, although a role in Th1 dominant diseases have been anticipated. Antagonists and gene knockouts have not been reported. In addition to T cell chemotaxis, CXCR3 ligands are angiostatic factors *in vivo*, but mechanisms are not defined.

C. CXCR4

CXCR4 is the first chemokine receptor shown to be an HIV-1 coreceptor (Feng et al., 1996) and the only one shown to be essential for life, at least in mice (Ma et al., 1998; Tachibana et al., 1998; Zou et al., 1998). Four groups identified it based on "orphan receptor" cloning strategies, whereas Feng et al. (1996) rediscovered the

cDNA by expression cloning of its HIV-1 coreceptor activity and named the protein "fusin." Specificity for the homeostatic CXC chemokine SDF-1 was established shortly thereafter (Bleul et al., 1996; Oberlin et al., 1996), and fusin was renamed CXCR4.

The ORF is interrupted by one intron in the region encoding the N-terminal segment and predicts a polypeptide 352 aa in length. A splice variant of unclear significance has been found, which affects the length of the N-terminal portion of the molecule upstream of TMD1, but not affinity for ligand (Heesen et al., 1997; Prodl et al., 1998; Gupta and Pillarisetti, 1999).

CXCR4 is unusually widely expressed on most hematopoietic cell types, including neutrophils, monocytes, T lymphocytes, B cells, B cell precursors, CD34⁺ progenitor cells from blood and bone marrow, blood-derived dendritic cells, Langerhans cells, T cells and macrophages, and both immature and mature T cells in thymus (Bleul et al., 1997; Zaitseva et al., 1997, 1998). It is also expressed at high levels on vascular endothelial cells (Gupta et al., 1998b), neurons from both the central and peripheral nervous systems (Hesselgesser et al., 1997), and microglia and astrocytes (He et al., 1997). In blood-derived T cells, CXCR4 is preferentially expressed on the naive, unactivated CD26low CD45RA⁺ CD45RO⁻ subset (Bleul et al., 1997), and expression is rapidly up-regulated by phytohemagglutinin and IL-2 (Loetscher et al., 1996) and down-regulated by SDF-1 (Amara et al., 1997).

CXCR4 has also been implicated in platelet formation. Although there is agreement over whether it is expressed throughout platelet development, there is some disagreement about its function (Power et al., 1995a; Hamada et al., 1998; Wang et al., 1998; Kowalska et al., 1999). SDF-1-induced transendothelial migration by mature marrow megakaryocytes and megakaryocyte progenitors has been reported by at least one group but not consistently confirmed. The receptor is on mature platelets but appears to be functionally uncoupled.

The SDF-1 gene is alternately spliced to form SDF-1 α and SDF-1 β , which differ by a 4-aa extension at the C terminus (Shirozu et al., 1995). These variants, originally isolated from bone marrow stromal cells, are functionally indistinguishable and are the only known endogenous ligands and agonists for CXCR4, inducing calcium flux and chemotaxis in transfected and primary cells in vitro. Genetic disruption of SDF-1 and CXCR4 in the mouse gives the same phenotype (Nagasawa et al., 1996; Ma et al., 1998; Tachibana et al., 1998; Zou et al., 1998), suggesting that they make up a monomeric signaling unit in vivo. The animals die in the perinatal period, the only known chemokine system components for which this is true, and have ventricular septal defects, defective gastric vasculogenesis and cerebellar development, abnormal bone marrow myelopoiesis, and defective B cell, but normal T cell, lymphopoiesis. Functions of CXCR4 in the adult are not defined. In one

study, human stem cell engraftment was reported to be regulated by CXCR4 in NOD/SCID mice (Peled et al., 1999). Both SDF-1 and CXCR4 have highly conserved sequences (e.g., 98 and 94% aa identity between human and mouse, respectively), which is highly atypical for chemokines and chemokine receptors, which are among the most rapidly evolving proteins in mammals (Murphy, 1993).

HIV-1 strains able to use CXCR4 for cell entry in vitro are named X4 strains (Berger et al., 1998). They are typically isolated late in the course of infection and correlate more or less with T cell line cytotropism and the syncytium-inducing methods of classification used before the discovery of HIV-1 coreceptors (reviewed in Berger et al., 1999). The importance of CXCR4 in HIV pathogenesis has been suggested but not proved by the detection of X4 HIV in CCR5-deficient HIV-positive individuals (Michael et al., 1998), and the discovery of a single nucleotide polymorphism in the 3'-UTR of SDF-1 α (SDF1-3'A) that is associated with slowed progression to AIDS (Winkler et al., 1998). Direct studies of the effect of this polymorphism on SDF-1 production in vivo have not been reported, but any effect could conceivably modulate the extent of X4 HIV interaction with CXCR4.

gp120 from HIV-1 envelope glycoprotein can bind to CXCR4 in the presence of CD4 (Lapham et al., 1999), and X4 virus entry is dependent on CD4 (Feng et al., 1996). However, CD4 independent association of gp120 to CXCR4 has also been demonstrated. Purified X4 gp120 can function as a CXCR4-dependent monocyte chemoattractant, perhaps to recruit more targets, and can induce apoptosis of the human neuronal cell line hNT (Hesselgesser et al., 1998a). Consistent with this, chemokines can regulate hippocampal neuronal signaling and gp120 neurotoxicity (Meucci et al., 1998). These findings may be relevant to the pathogenesis of HIV encephalitis and AIDS dementia. Interaction of gp120 with CXCR4 on macrophages can also induce apoptosis of CD8⁺ T cells, suggesting a coreceptor mechanism of CTL suppression (Herbein et al., 1998).

Several mAbs have been developed that bind CXCR4, including the prototype 12G5, which blocks HIV infection (Endres et al., 1996). Several small molecules and peptides, including some originally identified in HIV drug discovery programs, have been shown to selectively block chemokine receptor and/or HIV coreceptor activities of CXCR4. They include SDF-1 derived peptides (Loetscher et al., 1998b; Heveker et al., 1998); the synthetic peptide T22 (Tyr⁵,12,Lys⁷)polyphemus II, which consists of 18 aa residues and an analog of polyphemus II isolated from the hemocyte debris of American horseshoe crabs (*Limulus polyphemus*) (Murakami et al., 1997); the related synthetic peptides T134 and T140 (Tamamura et al., 1998; Xu et al., 1999); the polyarginine ALX40-4C (Doranz et al., 1997); the peptoid CGP64222 (Daelemans et al., 2000); and the bicy-

clam AMD3100 (Schols et al., 1997; Donzella et al., 1998; Bridger et al., 1999) (Fig. 3). The distamycin analog 2,2'-[4,4'-[aminocarbonyl]amino]bis[N,4'-di]pyrrole-2-carboxamide-1,1'-dimethyl]-6,8-naphthalenedisulfonic acid] hexasodium salt (NSC 651016) also blocks X4 viral use of CXCR4, but it has a broad specificity for multiple other chemokine receptors (Howard et al., 1998). CXCR4 has also been blocked with intrakines, which are modified forms of SDF-1 delivered by gene therapy that remain in the endoplasmic reticulum and block surface expression of newly synthesized CXCR4 (Chen et al., 1997).

In addition, the HIV protein Tat, which has a highly basic domain but lacks a chemokine fold, can block both SDF-1-induced calcium flux at CXCR4 and X4 HIV entry of target cells (Xiao et al., submitted). The inability of Tat to affect CCR5 function suggests a possible mechanism for restriction of HIV to R5 strains early in infection but cannot explain the appearance of X4 strains late in infection during immune system collapse. Moreover, it conflicts with the reported ability of Tat to up-regulate CXCR4 and serve as a vaccine target in nonhuman primates (Gallo, 1999).

The clinical development of CXCR4 blocking agents in HIV infection will have to confront safety questions of whether the virus will evolve to use other coreceptors and whether one or more of the phenotypes seen in CXCR4 knockout mice will occur. To date, CXCR4 has not been established as a therapeutic target for other diseases.

D. CXCR5

CXCR5 was the first chemokine receptor shown to be involved in lymphocyte homing and development of normal lymphoid tissue (Forster et al., 1996) and the first B cell selective chemokine receptor (Gunn et al., 1998a; Legler et al., 1998). Two cDNAs for CXCR5 were cloned independently by two groups as orphans and named, according to the source, monocyte-derived receptor 15 (MDR15; Barella et al., 1995) and Burkitt's lymphoma receptor 1 (BLR1; Dobner et al., 1992). The ORF of MDR15 has 327 codons and is 45 codons shorter at the N terminus than BLR1 due to alternative splicing of the gene. Distinct pharmacology has not been demonstrated for the two forms. The aa sequence is ~40% identical with CXCR1 and CXCR2.

Using a mAb directed to the N terminus of BLR1, CXCR5 has been detected on all peripheral blood and tonsillar B cells but only on a fraction of cord blood and bone marrow B cells. It is also present on a small subset of peripheral blood CD4⁺ (14%) and CD8⁺ (2%) T cells, which are also CD45RO⁺, IL-2R⁻, CD44^{high}, and L-selectin^{low}, suggesting a memory phenotype. In contrast, in secondary lymphatic tissue, the majority of CD4⁺ cells are positive, and in cord blood, T cells are negative (Forster et al., 1994). The murine homolog of CXCR5 has been cloned, and specific transcripts found in a pattern

similar to the human receptor, including expression on mature B cells and a subpopulation of T helper cells, as well as in secondary lymphatic organs and to a lesser extent in brain, specifically in the granule and Purkinje cell layer of the cerebellum (Kaiser et al., 1993). RNA in situ hybridization localizes transcripts to primary follicles and to the mantle zone of secondary follicles. Like other chemokine receptors, CXCR5 is dynamically regulated on T cells. After T cell receptor (TCR) stimulation, CXCR5 is up-regulated on memory/effector T cells, whereas IL-2 causes down-regulation (Sallusto et al., 1999b). Up-regulation of CXCR5 on antigen-activated T cells implies a role for movement of Th cells to B cell follicles (Ansel et al., 1999).

To date, B cell-attracting chemokine 1 (BCA-1, also known as BLC) is the only known agonist for CXCR5 (Gunn et al., 1998a; Legler et al., 1998). Conversely, CXCR5 is the only known receptor for BCA-1. Signaling includes chemotaxis and Ca²⁺ mobilization. BCA-1, a member of the homeostatic class of chemokines, is B cell selective and constitutively expressed in secondary lymphoid organs. It has weak effects on small numbers of T cells and macrophages. Consistent with this, CXCR5 knockout mice have a severe defect in normal B cell migration and localization (Forster et al., 1996). The animals lack inguinal lymph nodes, have few Peyer's patches, and have abnormal primary lymphoid follicles and no functional germinal centers in spleen. Nevertheless, immunoglobulin levels are normal. Disease phenotypes have not been reported.

Thus, although the biological importance of this receptor is established, evidence is lacking for its significance as a therapeutic target in disease. No CXCR5 antagonists or neutralizing mAbs have been developed yet. Recently, CXCR5 was reported to have coreceptor activity selective for HIV-2 (Kanbe et al., 1999).

IV. CC Chemokine Receptor Subtypes

A. CCR1

CCR1 was the first CC chemokine receptor identified and the first shown to have a functional viral homolog, US28 of human cytomegalovirus (Gao et al., 1993; Neote et al., 1993; Gao and Murphy, 1994). The gene is on human chromosome 3p21 in a cluster with CCR2, CCR3, CCR4, CCR5, CCR8, CCR9, XCR1, CX3CR1, and several orphans (Samson et al., 1996c). The ORF is on a single exon, and the predicted polypeptide is 355 aa in length.

Using a polyclonal rabbit antibody, Su et al. (1996) identified CCR1 on human peripheral blood lymphocytes and monocytes. A majority of CD3⁺, CD4⁺, CD8⁺, and CD16⁺ lymphocytes were positive. Among CD4⁺ peripheral blood T cells, CD45RO⁺ cells expressed greater amounts of CCR1 than CD45RO⁻ cells, suggesting selective expression on the memory subtype. Expression studies using an anti-CCR1 mAb have not been reported.

CCR1 binds multiple inflammatory/inducible CC chemokines with similar high affinity, including MIP-1 α , RANTES, MCP-2, MCP-3, leukotactin-1/MIP-5, MPIF-1 and HCC-1 (Neote et al., 1993; Gao et al., 1993; Youn et al., 1997; Gong et al., 1997; Tsou et al., 1998; Zhang et al., 1999; Nardelli et al., 1999). MIP-1 β and MCP-1 bind with much lower affinity and are poor agonists (Neote et al., 1993). HCC-1 may be selective. Mouse CCR1 (80% aa identity) binds human and mouse MIP-1 α with high affinity; agonists include mouse and human MIP-1 α and human leukotactin-1/MIP-5 (Gao and Murphy, 1995; Post et al., 1995; Zhang et al., 1999). A closely related mouse orphan named MIP-1 α -RL1 (65% aa identity) has also been cloned, but it has no human counterpart (Gao and Murphy, 1995).

CCR1 signaling includes calcium flux, inhibition of adenylyl cyclase, and chemotaxis (Myers et al., 1995; Pease et al., 1998). Coupling to both G $_i$ and G $_{14}$, but not G $_{11}$ or G $_{16}$, has been reported in transfected COS cells (Kuang et al., 1996). Signaling can be blocked efficiently by RANTES variants that have been modified at the N terminus, including truncated forms (Arenzana-Seisdedos et al., 1996; Struyi et al., 1998). Met-RANTES (Proudfoot et al., 1996), and amino-oxy-pentane (AOP)-RANTES (Simmons et al., 1997); however, none of these is selective for CCR1 over the other RANTES receptors, CCR3 and CCR5. High CCR1 selectivity has been reported by Berlex Biosciences for 4-hydroxypiperidines (K_i = 40–4000 nM) (Hesselgesser et al., 1998b; Ng et al., 1999) (Fig. 3), particularly 2,2-diphenyl-5-(4-chlorophenyl)piperidin-yl)valeronitrile, which inhibits MIP-1 α binding to CCR1 (K_i ~ 40 nM) and blocks MIP-1 α -induced extracellular acidification, Ca $^{2+}$ mobilization, and chemotaxis of peripheral blood mononuclear cells; effects in disease have not been reported yet. Other small molecule CCR1 antagonists have also been disclosed but have either lower potency or selectivity than the Berlex Biosciences compound.

Clear disease indications have not yet been identified for CCR1. Nevertheless, there is a fair amount now known about its biology from the phenotype of CCR1 knockout mice. The receptor is dispensable for growth, development, and reproduction, and the mice do not acquire spontaneous infections from environmental pathogens. It is the dominant receptor used by MIP-1 α for induction of mouse neutrophil chemotaxis and calcium flux in vitro, mobilization of neutrophils and hematopoietic progenitor cells in vivo, and regulation of hematopoietic progenitor cell proliferation (Gao et al., 1997; Broxmeyer et al., 1999). Consistent with this, MIP-1 α functions as a negative regulator of hematopoiesis (reviewed in Broxmeyer et al., 1997), and in vitro anti-CCR1 antibodies block MIP-1 α inhibition of colony formation by burst-forming unit-erythroid from purified human CD34 $^{+}$ bone marrow cells (Su et al., 1997). In this regard, BB10010, an agonistic variant of MIP-1 α (British Biotech, Inc.), has been tested in phase I and II

clinical trials as a stem cell protective agent in patients undergoing chemotherapy. The agent was safe in the doses tested, but only small therapeutic effects were noted on myeloipoiesis, perhaps because of insufficient stress on the bone marrow by the chemotherapy regimens tested (Clemons et al., 1998; Marshall et al., 1998). Another CCR1 agonist, MPIF-1 (Human Genome Sciences, Rockville, MD), has recently undergone phase I trial for the same indication.

Consistent with a role in neutrophils, CCR1 $-/-$ mice have reduced alveolitis in a pancreatitis-alveolitis mouse model (Gerard et al., 1997), as well as increased lethality when infected with *Aspergillus fumigatus*, an organism controlled primarily by neutrophils (Gao et al., 1997). However, this is an example where mouse and human orthologs may differ in biological function, because the major CCR1 agonists MIP-1 α and RANTES are poor agonists for human neutrophils (Coulain et al., 1997; Youn et al., 1997; Zhang et al., 1999). CCR1 also regulates granuloma formation and Th1/Th2 cytokine balance in response to Schistosoma eggs deposited in mouse lung, but it is not a dominant receptor for MIP-1 α -induced macrophage chemotaxis in vitro (Gao et al., 1997). Nevertheless, CCR1 deficiency did not reduce neutrophil accumulation in a nephrotic nephritis mouse model; disease was actually exacerbated with increased accumulation of macrophages and CD4 $^{+}$ and CD8 $^{+}$ T cells, as well as enhanced effector immune responses (Topham et al., 1999). However, CCR1 deficiency suppressed development of acute and chronic cardiac allograft rejection in several mouse models (Gao et al., 2000). Thus CCR1 can modulate inflammatory responses either positively or negatively, depending on the context, through effects on multiple leukocyte subtypes. The phenotype of MIP-1 α knockout mice includes protection from coxsackievirus myocarditis, influenza A alveolitis, and acute experimental allergic encephalomyelitis; however, specific roles for CCR1 are not defined (Cook et al., 1995; Kennedy et al., 1998).

B. CCR2

CCR2 is the only leukocyte MCP-1 receptor identified so far, and it is important in inflammation, including atherosclerosis. The ORF is on two alternatively spliced exons that encode two distinct polypeptides 360 (CCR2 $_{(a)}$) and 374 (CCR2 $_{(b)}$) aa in length (Charo et al., 1994; Wong et al., 1997). The two have an identical sequence until aa 313, which is located in the C-terminal cytoplasmic region, and similar functional properties. Both RNAs are detectable in monocytes, blood-derived DC and NK cells and T lymphocytes but not in resting neutrophils or eosinophils. CCR2 $_{(b)}$ appears to be the predominant form. mAbs have identified functional CCR2 in monocytes, activated memory T cells, B cells, and basophils (Frade et al., 1997; Rabin et al., 1999). In vivo, chronic inflammation may potentiate neutrophil migration to MCP-1 (Johnston et al., 1999).

Signaling through CCR2 in transfected cells includes calcium mobilization, inhibition of adenyllyl cyclase, and chemotaxis (Myers et al., 1995). Receptor triggering may require receptor dimerization (Rodriguez-Frade et al., 1999). CCR2 binds multiple inflammatory/inducible ligands with similar high affinity, including MCP-1, MCP-2, MCP-3, MCP-4, and mouse MCP-5 (Charo et al., 1994; Ben-Baruch et al., 1995; Garcia-Zepeda et al., 1996; Gong et al., 1997; Sarafi et al., 1997). Only MCP-1 is selective versus other chemokine receptors, although it also binds to D6 and Duffy (see later). The HIV Tat protein is also an agonist at CCR2, which has suggested a possible mechanism for recruitment of target cells to sites of HIV infection (Albini et al., 1998).

mAb MCP-1 R02 directed to the CCR2 N terminus is also an agonist, whereas mAbs directed to the third extracellular domain (MCP-1R04 and MCP-1 R05) are antagonists (Rodriguez-Frade et al., 1999). Small molecule CCR2 antagonists have been reported in the patent literature by Roche Biosciences (Fig. 3).

Mouse CCR2 has 80% aa identity to human CCR2, is expressed in peritoneal macrophages, and is specific for the mouse chemokines JE and FIC, which have highest sequence homology to MCP-1 and MCP-3, respectively (Boring et al., 1996; Kurihara and Bravo, 1996). Using chemokine neutralization in mice, Karpus's group found that acute and relapsing forms of experimental autoimmune encephalomyelitis are regulated by differential expression of MIP-1 α and JE/MCP-1, respectively, implicating CCR2 in relapsing forms of EAE (Kennedy et al., 1998). Correlative studies have also implicated chemokines and chemokine receptors in the pathogenesis of multiple sclerosis (Ransohoff, 1999).

Mice lacking CCR2 develop normally but do not recruit macrophages in an experimental peritoneal inflammation model, fail to clear *Listeria monocytogenes*, have smaller granulomas after i.v. injection with yeast β -glucan, and have smaller granulomas and lower production of interferon- γ in draining lymph nodes when challenged with immobilized PPD by emulsification to lung (Boring et al., 1997; Kurihara et al., 1997; Kuziel et al., 1997). As for CCR1 knockouts, this suggests a role in immunomodulation as well as in direct recruitment of monocytes/macrophages to sites of inflammation. They also have defective cockroach allergen-induced bronchial hyperreactivity (Campbell et al., 1999). Consistent with a pathogenetic role for MCP-1 and macrophages in human atherosclerotic plaques, CCR2 $-/-$ mice have a sustained ~50% reduction in size of atherosclerotic lesions when challenged with a Western diet on an apolipoprotein E $-/-$ genetic background, which normally produces severe atherosclerosis (Boring et al., 1998). This is consistent with results from similar studies of JE/low-density lipoprotein receptor double knockout mice and JE deficiency in mice overexpressing human apolipoprotein B (Gu et al., 1998; Gosling et al., 1999). The protective effects are not mediated by changes in

lipid levels and occur at both high and low levels of plasma lipids. The effect of blocking CCR2 in established atherosclerosis has not been reported yet.

CCR2_b has HIV-1 coreceptor activity in vitro, but the activity and strain specificity are both low (compared with CCR5 and CXCR4; Doranz et al., 1996; Zhang et al., 1998a). Nevertheless, a role in disease has been suggested by discovery of a common variant CCR2 allele named CCR2-64I that is associated with a 2- to 4-year delay in progression to AIDS in some HIV-1 seroconverter cohorts (Smith et al., 1997). To date, the mechanism of action of this mutation has not been defined. In particular, it does not appear to directly affect either the chemokine receptor or HIV coreceptor activities of CCR2 or other coreceptors (Lee et al., 1998). Interestingly, CCR2-64I but not wild-type receptor has been observed to heterodimerize with native CXCR4 when transfected in human embryonic kidney 293 cells; however, functional correlates have not been defined (Mellado et al., 1999). CCR2-64I could also potentially be linked to a disease-modifying mutation in CCR5, because the two genes are located within 10 kb on chromosome 3.

C. CCR3

CCR3 is an eosinophil chemoattractant receptor for multiple inflammatory/inducible CC chemokines (Heath et al., 1997) and may be important in allergic inflammation, including asthma, and antihelminthic host defense where eosinophils greatly outnumber other leukocytes; it is also an HIV-1 coreceptor (Doranz et al., 1996; Choe et al., 1996). A human CCR3 cDNA highly expressed in eosinophils was first reported by Combadiere et al. (1995a). Its ligands were originally reported incorrectly as MIP-1 α , MIP-1 β , and RANTES and later corrected to eotaxin (Kitaura et al., 1996). Two other groups independently characterized CCR3 as an eotaxin receptor and identified additional agonists (Daugherty et al., 1996; Ponath et al., 1996). The ORF is on a single exon and predicts a polypeptide 355 aa in length. CCR3 is most similar to CCR1 in sequence (62% aa identity) and ligands.

CCR3 is difficult to express in foreign cells, and ligand binding is highly dependent on pH and salt concentration (Dairaghi et al., 1997). Ligands and agonists for human CCR3 include eotaxin, eotaxin-2, eotaxin-3, RANTES, MCP-3, MCP-4, MIP-5/leukotactin-1, and HIV Tat (Daugherty et al., 1996; Garcia-Zepeda et al., 1996; Kitaura et al., 1996, 1999; Ponath et al., 1996; Forssmann et al., 1997; Gong et al., 1997; Heath et al., 1997; Youn et al., 1997; Albini et al., 1998; Shinkai et al., 1999). Eotaxin, eotaxin-2, and eotaxin-3 appear to be the most potent and are selective.

Human CCR3 distribution on eosinophils (Heath et al., 1997), basophils (Uguccioni et al., 1997), mast cells (Ochi et al., 1999), and a subset of Th2 T lymphocytes (Gerber et al., 1997; Sallusto et al., 1997) is compatible with a role in allergic inflammation. The receptor is also

found on dendritic cells (Rubbert et al., 1998) and microglial cells of the brain (He et al., 1997). CCR3 activities on human cells *in vitro* include eosinophil arrest under flow conditions (Kitayama et al., 1998), eosinophil and Th2 cell chemotaxis (Heath et al., 1997; Sallusto et al., 1997), degranulation of eosinophils and basophils (Uguccioni et al., 1997), HIV-1 entry of microglial cells (He et al., 1996), and HIV-specific T cell cytotoxicity mediated by RANTES (Hadida et al., 1998).

A fully antagonistic mAb specific for human CCR3 named 7B11 was developed and used to show that eosinophil and basophil responses to eotaxin, RANTES, MCP-2, MCP-3, and MCP-4 are mediated entirely by CCR3 in most donors (Heath et al., 1997; Uguccioni et al., 1997). The chemokine derivative Met-chemokine $\beta 7$ is a potent and specific antagonist (Nibbs et al., 2000). Less specific antagonists include Met-RANTES, AOP-RANTES, the distamycin analog previously mentioned in the section on CXCR4, and vMIP-II (Kledal et al., 1997). It has also been reported that vMIP-II chemottracts human eosinophils in a CCR3-dependent manner, a discrepancy that has not been reconciled (Boshoff et al., 1997). Most of these blocking agents also block HIV interaction with CCR3. A small molecule antagonist has been reported by Takeda in the patent literature (Fig. 3).

The discovery of eotaxin and CCR3 raised hopes that this would be a dominant signaling system in allergic inflammation and a major new drug target; however, so far, the evidence from animal models is inconclusive. Eotaxin knockout mice have been generated by two groups, but reported effects on airway eosinophilia after ovalbumin challenge have conflicted: Rothenberg et al. (1997) reported a ~40% reduction, whereas Yang et al. (1998) saw no effect. Eotaxin is required for the baseline level of tissue eosinophils (Matthews et al., 1998). A CCR3 knockout mouse has not yet been reported. Mouse may be a poor host for modeling eotaxin and CCR3, because CCR1 is highly expressed in mouse versus human eosinophils (Gao et al., 1996); mouse CCR3, unlike human CCR3, binds MIP-1 α in addition to eotaxin (Post et al., 1995); and mouse CCR3 is expressed only on eosinophils (Grimaldi et al., 1999). The guinea pig has been proposed as a superior model, which can now be studied with a neutralizing mAb recently developed against guinea pig CCR3 (Sabroe et al., 1998). Meanwhile, previous work on the neutralization of guinea pig eotaxin revealed partial blockade of airway eosinophilia on allergen challenge (Humbles et al., 1997) and partial suppression of eosinophil mobilization from bone marrow (Palframan et al., 1998).

CCR3 has broad specificity for R5-, X4-, and dual-tropic HIV envelope glycoproteins and is used by HIV for entry of microglial cells *in vitro* (Choe et al., 1996; He et al., 1997; Bazan et al., 1998). However, use *in vivo* is not established.

D. CCR4

CCR4 has been reported to be a selective marker for Th2 T lymphocytes and is up-regulated by T cell receptor activation (Bonecchi et al., 1998; Sallusto et al., 1998, 1999b). Current concepts of CCR4 function include dendritic cell trafficking, T cell recirculation from tissue to draining lymph node, T cell transmigration through thymus during T cell maturation, T cell migration to ectopic lymphoid tissue (Sozzani et al., 1999), and homing of memory T cells to inflamed skin but not to gut (Campbell et al., 1999).

The cDNA was originally cloned from a human basophilic leukemia cell line library (Power et al., 1995). The ORF is on a single exon and predicts a polypeptide 360 aa in length. High-affinity ligands and high-potency agonists include MDC and TARC (Imai et al., 1997a, 1998), which are constitutively made and selective for CCR4. Activities include calcium flux and chemotaxis. MIP-1 α , RANTES, and MCP-1 are also agonists when CCR4 is expressed in frog oocytes and transmembrane currents are measured, but potencies have not been reported (Power et al., 1995). A cDNA encoding a mouse counterpart of CCR4 has been isolated that has 80% aa identity (Hoogewerf et al., 1996). Knockout mice, neutralizing mAbs, and small molecule antagonists have not been reported yet.

At present, potential roles for CCR4 in disease have been inferred based on analysis of its known ligands in mouse disease models. In one study, it was implicated in the pathogenesis of bacteria-induced fulminant hepatic failure in mice, based on a protective effect of injecting anti-TARC mAbs (Yoneyama et al., 1998). In a second study, neutralization of mouse MDC was protective in a model of airway hyperreactivity and eosinophilic inflammation (Gonzalo et al., 1999).

Purified native human MDC has been reported to be a broad-spectrum HIV-1 suppressive agent *in vitro*, but its mechanism of action is unlikely to involve CCR4, because it has not been observed to function as an HIV coreceptor and because TARC lacks this activity (Pal et al., 1997). Several laboratories have been unable to reproduce this effect with recombinant and synthetic MDC.

E. CCR5

CCR5 is a major HIV-1 coreceptor that controls susceptibility to HIV-1 infection and disease. The first report of the sequence and ligands for human CCR5 was published on March 19, 1996 (Samson et al., 1996a), followed on March 29 and June 14 by mouse CCR5 (Boring et al., 1996; Meyer et al., 1996) and in July 1996 by two additional independent reports of human CCR5 (Combadiere et al., 1996; Raport et al., 1996). The HIV coreceptor activity was described in five reports by five independent groups published within 1 week in June 1996 (Alkhatib et al., 1996; Choe et al., 1996; Deng et al.,

1996; Dragic et al., 1996; Doranz et al., 1996), and 2 months later the first of a series of reports describing the defective CCR5 Δ 32 allele, which established the pivotal *in vivo* role of CCR5 in HIV pathogenesis, was reported (Dean et al., 1996; Huang et al., 1996; Liu et al., 1996; Samson et al., 1996b; Zimmerman et al., 1997).

The human CCR5 ORF is on a single exon and predicts a polypeptide 355 aa in length. It is expressed on peripheral blood-derived dendritic cells (Granelli-Piperno et al., 1996; Rubbert et al., 1998), CD34⁺ hematopoietic progenitor cells (Ruiz et al., 1998), and activated/memory CD26^{high} CD45RA^{low} CD45RO⁺ Th1 lymphocytes (Bleul et al., 1997; Loetscher et al., 1998c). Fresh monocytes express low levels that can be increased by culture *in vitro* (Alkhatib et al., 1996). Likewise, freshly isolated T cells express low amounts that increase with prolonged stimulation by IL-2, mitogens, and other activating protocols *ex vivo* (Bleul et al., 1997) or by Th1 type inflammation *in vivo* (e.g., in synovial fluid from patients with rheumatoid arthritis; Qin et al., 1998). CCR5 is also expressed on CD4⁺ and CD8⁺ thymocytes (Zaitseva et al., 1998) and Langerhans cells (Zaitseva et al., 1997). Reports of CCR5 on neurons, astrocytes, capillary endothelial cells, epithelium, vascular smooth muscle, and fibroblasts have also been published, but functional roles are not established.

In vivo, T lymphocytes and macrophages in both lymphoid and nonlymphoid tissues express CCR5 and CXCR4, but follicular dendritic cells in lymph node express neither, suggesting that trapping of HIV by these cells does not involve the major HIV coreceptors. CCR5-positive cells are more frequently identified in the colon than in the rectum and more frequently identified in the cervix than in the vagina, suggesting that the expression levels of coreceptors are differentially regulated at different anatomic sites (Zhang et al., 1998b).

High-affinity ligands and high-potency agonists include MIP-1 α (P form is more potent than S form; Nibbs et al., 1999), RANTES, MIP-1 β , and MCP-2, but none is selective (Combadiere et al., 1996; Raport et al., 1996; Samson et al., 1996a; Gong et al., 1998). Additional ligands include MCP-3, MCP-4, MCP-1, and eotaxin. MCP-3 appears to be an antagonist (Blanpain et al., 1999). gp120 from R5 HIV strains are also ligands and induce receptor triggering and chemotaxis in a CD4-dependent manner (Weissman et al., 1997). CCR5 has been found to associate constitutively with CD4 (Xiao et al., 1999), but the physiological role is unknown. Chemokine ligands block CCR5 use by R5 HIV-1 strains, and vice versa (Cocchi et al., 1995; Alkhatib et al., 1996; Trkola et al., 1996).

CCR5 blocking agents include mAbs, some of which selectively block HIV coreceptor activity but not chemokine binding (Wu et al., 1997; Olson et al., 1999), and chemokine derivatives, such as truncated versions of RANTES, Met-RANTES, and AOP-RANTES and the viral chemokine KSHV vMIP-II, all of which block both

chemokine and HIV interaction with CCR5 but are not selective (Arenzana-Seisdedos, 1996; Kledal et al., 1997; Simmons et al., 1997). Met-RANTES and AOP-RANTES are pure and partial antagonists at CCR5, respectively. AOP-RANTES induces calcium flux but not chemotaxis and is the most potent RANTES derivative in blocking R5 HIV entry. Moreover, it blocks HIV entry of all target cells tested, whereas wild-type RANTES, for unclear reasons, fails to efficiently block R5 HIV entry of macrophages.

CCR5 can be blocked selectively by mAbs; by intrakines, which are chemokines delivered by gene therapy and targeted for endoplasmic reticulum retention that trap CCR5 intracellularly (Yang et al., 1997); by hammerhead ribozymes (Goila and Banerjee, 1998); and by the small molecule TAK-779 (*N,N*-dimethyl-*N*-[4-[[[2-(4-methylphenyl)-6,7-dihydro-5*H*-benzocyclohepten-8-yl]carbonyl]amino]benzyl]tetrahydro-2*H*-pyran-4-aminium chloride) (Baba et al., 1999) (Fig. 3). Moreover, a novel fusion-competent HIV vaccine strategy has been discovered in which CCR5, CD4, and a cross-linking agent are used to trap neutralizing epitopes of gp120 for presentation to the immune system (LaCasse et al., 1999). Immunization of mice led to the production of neutralizing antisera for diverse primary HIV isolates.

CCR5 is blocked naturally by inheritance of CCR5 Δ 32, a mutant allele common in whites that encodes a truncated, inactive receptor due to a 32-bp deletion in the ORF (Dean et al., 1996; Huang et al., 1996; Liu et al., 1996; Samson et al., 1996b; Zimmerman et al., 1997). CCR5 Δ 32 homozygotes, which represent ~1% of North American whites, exhibit high resistance to HIV infection and appear otherwise healthy. This suggests that normal CCR5 function is well compensated or redundant. CCR5 Δ 32 heterozygotes have reduced normal CCR5 expression on cells, due in part to a dominant negative effect of the CCR5 Δ 32 protein (Benkirane et al., 1997) and, if infected with HIV, progress less rapidly to AIDS in some but not all cohorts that have been studied. An even stronger effect on disease progression has been associated with a single nucleotide polymorphism located in the CCR5 promoter (P1 or 59029 G/A alleles), which affects gene transcription (McDermott et al., 1998; Martin et al., 1998). Why the CCR5 Δ 32 allele is so common in whites is unknown, but a reasonable mechanism is selection through an earlier epidemic within the past two millennia (Libert et al., 1998). In this regard, it is interesting to note that myxoma, a rabbit poxvirus, can use CCR5 (as well as several other chemokine receptors) as a cell entry factor (Lalani et al., 1999). Perhaps variola could also use CCR5 but not CCR5 Δ 32, which was enriched in populations by smallpox epidemics.

Mouse CCR5 is similar in ligand selectivity to the human receptor, and CCR5 $-/-$ mice, like CCR5-deficient humans, appear healthy (Zhou et al., 1998). Subtle defects have been identified in stressed mice, including

reduced efficiency in the clearance of *Listeria* infection, relative resistance to lipopolysaccharide-induced endotoxemia, increased susceptibility to *Cryptococcus* infection (Huffnagle et al., 1999), enhanced delayed-type hypersensitivity reaction, and increased humoral responses to T cell-dependent antigenic challenge, indicating a role for CCR5 in down-modulating T cell-dependent immune responses.

To date, CCR5 is the only chemokine receptor (excluding Duffy for the moment) for which proof of concept is available for a role in human disease. The development of therapeutic and preventive strategies that mimic the near-perfect safety and efficacy of CCR5Δ32 in the prevention of infection of HIV-exposed populations is therefore of substantial interest. However, blocking CCR5 alone in the setting of established infection is unlikely to be effective because viruses are likely to emerge that can use CXCR4 or other coreceptors. To date, one phase I clinical trial has addressed this question, using the MIP-1α variant BB10010 as a CCR5 blocking agent in HIV-positive individuals. No significant effect was observed on viral burden or CD4 counts, but the maximal administered dose was insufficient to achieve blocking levels in vivo (L. Czaplewski, personal communication).

F. CCR6

CCR6 is the only known receptor for LARC (also known as MIP-3α, exodus and ck-84) (Baba et al., 1997; Greaves et al., 1997; Liao et al., 1997b; Power et al., 1997), and it mediates responsiveness of diverse subsets of memory T cells to LARC (Liao et al., 1999). Additional unusual features include its location on human chromosome 6q27 outside of the main CCR cluster on 3p (Liao et al., 1997a), its functional expression on nonactivated memory T cells (Campbell et al., 1998; Liao et al., 1999), and down-regulation during dendritic cell maturation (Dieu et al., 1998). Although the biology and pharmacology of CCR6 are not yet developed, it is predicted to be important in memory T cell and dendritic cell trafficking to secondary lymphoid organs. The CCR6 ligand MIP-3α appears to specifically regulate constitutive homing of Langerhans-type dendritic cells to the epidermis, whereas other types of dendritic cells respond to multiple other chemokines (Charbonnier et al., 1999).

The ORF is on two exons and predicts a polypeptide 368 or 374 aa in length, depending on which of two deduced methionines is considered to initiate translation. The aa sequence is 76% identical with mouse CCR6 (Varona et al., 1998). CCR6 mRNA is present constitutively in secondary lymphoid tissue (spleen, lymph nodes, appendix) and fetal liver, in peripheral blood CD4⁺ and CD8⁺ T cells with a memory phenotype, and in B cells but not NK cells, monocytes, or granulocytes. It is also selectively expressed in human dendritic cells derived from CD34⁺ cord blood precursors and in dendritic cells derived from peripheral blood monocytes

(Yang et al., 1999a). TCR activation of T cells causes down-regulation of CCR6 (Sallusto et al., 1999b).

LARC, which is produced by activated macrophages, dendritic cells, and endothelial cells, is the only high-affinity ligand ($K_d = 0.1\text{--}12\text{ nM}$ on transfected cells, 0.4 nM on lymphocytes) and high-potency chemokine agonist for CCR6. Signaling includes calcium flux and chemotaxis. Recently, the human β -defensin HBD2 has been identified as a nonchemokine functional ligand for CCR6. HBD2 is produced by epithelial cells during infection and functions as a direct antimicrobial factor but also chemoattracts CCR6⁺ dendritic cells and memory T cells, suggesting a chemokine receptor link between innate and adaptive immunity (Yang et al., 1999b). The activity of other defensin family members at chemokine receptors has not been reported.

G. CCR7

CCR7 is a major homing receptor of the immune system, critical not only for trafficking of B lymphocytes, T lymphocytes, and dendritic cells across high endothelial venules but also for their correct positioning in T cell zones of secondary lymphoid organs (Cyster et al., 1999; Forster et al., 1999; Sozzani et al., 1999). A favored model views CCR7 as a homing switch that is turned on during the activation of resting T cells and maturing dendritic cells (Gunn et al., 1998b; Sallusto et al., 1998, 1999b; Sozzani et al., 1998a; Yanagihara et al., 1998; Kellerman et al., 1999; Forster et al., 1999; Saeki et al., 1999). Its importance is revealed by the profound disorganization of secondary lymphoid tissue in CCR7 $^{-/-}$ mice and the failure of these mice to mount rapid antibody responses as well as contact sensitivity and delayed-type hypersensitivity responses to T-dependent antigens (Forster et al., 1999).

CCR7 cDNAs were cloned first from B cells by two groups working independently: one using an Epstein-Barr virus-infected versus uninfected B lymphocyte subtraction cDNA library (Birkenbach et al., 1993), and the other using a Burkitt's lymphoma cell library. Thus, the receptor was originally named EB1-1, for Epstein-Barr virus-induced cDNA #1, and BLR-2, for Burkitt's lymphoma receptor-2 (Burgstahler et al., 1995). The gene is on human chromosome 17q12-21.2 outside the main CCR cluster on 3p21 (Schweickart et al., 1994). The ORF is on two exons separated by an intron in the N-terminal domain and predicts a polypeptide 378 aa in length.

ELC and SLC are highly specific functional ligands for CCR7 that bind with similar affinity and induce chemotaxis of CCR7-positive cells (Yoshida et al., 1997, 1998a; Campbell et al., 1998). Both ELC and SLC are expressed constitutively within the T cell zone of secondary lymphoid tissue and mucosa-associated lymphoid tissue but not in B cell areas or sinuses or blood-derived leukocytes (Willmann et al., 1998). SLC expression has been finely mapped to interdigitating dendritic cells and high endothelial venules of secondary lymphoid tissue and in lymph

phatic endothelium of various organs, where it promotes adhesion and chemotaxis of naive T lymphocytes (Gunn et al., 1998a). This is consistent with the phenotype of mice homozygous for the spontaneous mutation *plt* (paucity of lymph node T cells), which maps to the SLC locus and in which SLC is not produced. The exact *plt* mutation has not been identified, but it is not within SLC exons or introns (Gunn et al., 1999). In these mice, naive T cells fail to home to lymph nodes or lymphoid regions of spleen, and skin dendritic cells fail to traffic to and accumulate in spleen and lymph node T cell zones. The defect is associated with markedly increased sensitivity to infection with murine hepatitis virus. The phenotypes of CCR7 $-/-$ mice and *plt* mice are very similar (Forster et al., 1999). Defective B and T cell homing in CCR7 $-/-$ mice has been shown by adoptive transfer experiments to be due to defective cell migration.

CCR7 is also dynamically regulated during thymocyte maturation; however, a defect in T cell development was not found in CCR7 $-/-$ mice (Forster et al., 1999; Suzuki et al., 1999). Evidence for chemoattraction of activated T lymphocytes by maturing dendritic cells via up-regulation of CCR7 ligands has also been reported (Tang and Cyster, 1999).

Using an anti-CCR7 mAb, CCR7 phenotype has been used to define subsets of T cells that mediate two functionally distinct aspects of immunological memory (Salustio et al., 1999). CCR7 negative memory cells, designated as TEM (effector memory T cells), express receptors for migration to inflamed tissue and have immediate effector function. CCR7+ memory cells or T_{CM} (central memory T cells) express lymph node homing receptors, efficiently stimulate dendritic cells, and differentiate into TEM on secondary stimulation but lack immediate effector function.

H. CCR8

CCR8 is notable for high expression in thymus (Napolitano et al., 1996) and association with Th2 lymphocytes (Zingoni et al., 1998). It was molecularly defined as a previously cloned orphan receptor (Roos et al., 1997; Tiffany et al., 1997; Goya et al., 1998; Horuk et al., 1998). The ORF is on a single exon and predicts a polypeptide 355 aa in length, 32 to 45% identical with other CCR subtypes. Mouse CCR8 has 71% aa identity to human CCR8 (Goya et al., 1998).

In addition to thymus and Th2 lymphocytes, CCR8 mRNA is found in brain, spleen, lymph node, and monocytes (Napolitano et al., 1996; Tiffany et al., 1997). I-309 is a selective human ligand; however, interestingly, it also binds three viral chemokines: vMIP-I and vMIP-II from KSHV and MC148R (also called vMCC-I) from *M. contagiosum* virus (Damon et al., 1998; Dairaghi et al., 1999; Endres et al., 1999). In calcium flux and chemotaxis assays, I-309 and vMIP-I are agonists, whereas MC148R is an antagonist; vMIP-II has been reported as an antagonist by one group (Dairaghi et al., 1999) and as

a chemotactic agonist by another (Sozzani et al., 1998b). TARC and MIP-1 β have also been reported as chemotactic agonists (Bernardini et al., 1998), but this has not been confirmed (Dairaghi et al., 1999; H. L. Tiffany and P. M. Murphy, unpublished data). Mouse CCR8 is also expressed in thymus. Both I-309 and its mouse homolog TCA-3 are high-affinity ligands and potent agonists at mouse CCR8 (Goya et al., 1998).

The biological roles played by CCR8 and its ligands are speculative at this point. Knockout mice, mAbs, and antagonists are currently unavailable. Expression of CCR8 in Th2 cells correlates well with the ability of I-309 to chemoattract these cells and suggests a possible role in allergic inflammation. Selective action of viral chemokines at CCR8 suggests a role in Kaposi's sarcoma (KS) and *M. contagiosum*, possibly through the modulation of Th2 function. Of note, Th2 cells are found in KS lesions, whereas *M. contagiosum* lesions are notable for the paucity of leukocytes. Apart from migration, the ability of I-309 to inhibit apoptosis of dexamethasone-treated mouse thymic lymphoma cells has suggested that CCR8 may be involved in this and possibly other apoptotic processes (Van Snick et al., 1996).

In transfected cells, CCR8 can function as an HIV-1 coreceptor for diverse T-cell tropic, dual-tropic, neurotropic, and macrophage-tropic HIV-1 strains, and I-309 can inhibit this activity (Horuk et al., 1998). However, the use of CCR8 by HIV-1 in primary cells and in pathogenesis is undefined.

I. CCR9

CCR9 is the previously cloned orphan GPR 9-6 (Youn et al., 1999; Yu et al., 2000; Zaballos et al., 1999; Zabel et al., 1999). The chemokine binding protein D6 had previously been inappropriately named CCR9 (see later). Two splice variants, designated CCR9_(a) and CCR9_(b), have been identified. CCR9_(a) is predicted to have 12 additional aa at its N terminus (compared with CCR9_(b)) and is the predominant form in all cell types examined (Yu et al., 2000). TECK, a homeostatic/constitutive type chemoattractant for dendritic cells, thymocytes, intestinal homing T lymphocytes, mucosal lymphocytes, and activated macrophages, is the only known agonist, and it acts with slightly higher potency at CCR9_(a) than CCR9_(b). Constitutive expression of human and mouse CCR9 is very high in thymus and low in lymph nodes and spleen. Immature and mature thymic T cells express CCR9, suggesting a role in T cell development in the thymus.

J. CCR10

CCR10 is the former orphan receptor GPR2 (Marchese et al., 1995). The only ligand identified to date is the placenta and skin-associated CC chemokine CCL27 (also known as ESkin, skinkine, and CTACK), which attracts skin-homing memory T cells (Baird et al., 1999;

Morales et al., 1999; Homey et al., 2000; C. Gerard et al., manuscript submitted).

K. CCR11

CCR11 is the second MCP-1 receptor identified. However, it has not yet been detected on leukocytes. The gene maps to 3p22 separate from the major CCR cluster at 3p21. Chemotactic ligands include MCP-1, MCP-2, and MCP-4, and it is expressed in heart, small intestine, and lung (Schweickart et al., 2000).

V. CX3C Chemokine Receptor Subtypes

A. CX3CR1

CX3CR1 is unique among chemokine receptors in functioning directly as a cell-cell adhesion, including under physiological shear (Imai et al., 1997b; Fong et al., 1998; Haskell et al., 1999). Its powerful adhesive action may be important for leukocyte extravasation in the setting of high blood flow. CX3CR1 has highest similarity to CC chemokine receptors (30–42%); consistent with this, the gene is located on human chromosome 3p21 in the major CCR cluster (Combadiere et al., 1995b).

As revealed first in rat and later in human and mouse, CX3CR1 mRNA is expressed at highest levels in brain, but it is found in all organs examined (Harrison et al., 1994; Combadiere et al., 1995b, 1998a). The only known ligand is fractalkine, which is also referred to as neurotactin in the mouse, a multimodular protein containing a chemokine domain, a mucin-like stalk, a transmembrane domain, and a cytoplasmic domain, as well as the CX3C signature (Bazan et al., 1997; Imai et al., 1997b; Pan et al., 1997). Fractalkine is found in a 95-kDa shed form as well as a membrane-tethered form expressed on endothelial cells and neurons (Bazan et al., 1997; Imai et al., 1997b; Pan et al., 1997; Harrison et al., 1998). Chemotactic signaling by CX3CR1 is G protein-mediated (pertussis toxin-sensitive), whereas direct adhesion to tethered fractalkine is not (Imai et al., 1997b). Blood-derived neutrophils, monocytes, NK cells, and T lymphocytes are positive for CX3CR1 and respond chemotactically to fractalkine (Bazan et al., 1997; Imai et al., 1997b). The recombinant chemokine domain of neurotactin is chemotactic for neutrophils both in vitro and in vivo (Pan et al., 1997). Membrane-bound fractalkine, in addition to being proadhesive, can also trigger receptors to signal and induces chemotaxis.

CX3CR1 appears to mediate cell adhesion to both endothelial cells and neurons. In rat brain, fractalkine has been detected on neurons, where it is up-regulated by axonal injury and mediates direct interactions with CX3CR1-expressing microglia, possibly to mediate nerve repair (Harrison et al., 1998). In the Wistar-Kyoto rat crescentic glomerulonephritis model, a setting of high blood flow, fractalkine is markedly induced in glomerular endothelium and CX3CR1 is increased on infil-

trating leukocytes. Moreover, anti-CX3CR1 antibody treatment dramatically inhibits leukocyte infiltration in the glomeruli, prevents crescent formation, and improves renal function (Feng et al., 1999). Effects on established disease are not defined.

In transfected cells, CX3CR1 can function as an HIV-1 coreceptor for a limited number of HIV-1 strains, and fractalkine can block this activity (Combadiere et al., 1998b). However, the use of CX3CR1 in primary cells by HIV-1 and in pathogenesis is undefined.

VI. Chemokine Receptor Subtypes

A. XCR1

XCR1 is the former orphan receptor GPR5 and the only C chemokine receptor, specific for the T lymphocyte directed molecule lymphotactin (Yoshida et al., 1998b). Biology and pharmacology have not yet been developed. However, a potential role in cancer therapy was suggested by the ability of lymphotactin to synergize with IL-2 in producing an antitumor immune response and tumor regression in mice (Dilloo et al., 1996).

VII. Chemokine Binding Proteins

A. Duffy

Duffy is the red cell receptor for the malaria-causing protozoan *Plasmodium vivax* and a highly promiscuous chemokine binding protein, specific for some but not all CC and CXC chemokines, such as ELR+ (IL-8, GRO α) but not ELR- CXC chemokines and basic (RANTES, MCP-1) but not acidic (MIP-1 α) CC chemokines (Miller et al., 1975, 1976; Darbonne et al., 1991; Chaudhuri et al., 1993, 1994; Horuk et al., 1993). Despite having a 7TMD architecture (338 aa) and ~25% aa identity to chemokine receptors, no signaling function has been observed on chemokine ligation (Neote et al., 1994). Because of its long history in the malaria and blood group literature (Horuk, 1994), the nomenclature committee has decided that the name of this molecule may remain Duffy even if signaling is demonstrated at some future date.

Normally, Duffy RNA is present in multiple organs, including brain, spleen, bone marrow, and lung, and in addition to red cells, Duffy antigen can be detected on endothelial cells of postcapillary venules (Hadley et al., 1994; Chaudhuri et al., 1997) and Purkinje cells of the cerebellum (Horuk et al., 1997). However, Duffy is selectively missing from red cells of individuals from equatorial and southern Africa (Chaudhuri et al., 1995; Peiper et al., 1995), the result of fixation of a variant allele bearing a single nucleotide mutation named -46C in an erythroid-specific GATA-1 site in the Duffy promoter (Tournamille et al., 1995). Accordingly, red cells from these individuals do not bind chemokine or plasmodium ligands for Duffy (Horuk et al., 1993; Chitnis and Miller, 1994); the individuals are resistant to *P.*

vivax; and *P. vivax* is virtually absent from this region of Africa. Analogous to CCR5 and HIV, these individuals have no obvious health problems attributable specifically to Duffy deficiency on red cells. Furthermore, an apparently healthy white individual has been described with complete Duffy deficiency, the result of a 14-bp deletion in the ORF (Mallinson et al., 1995). Together these experiments of nature suggest that Duffy is a dispensable gene. This presents a challenge to hypotheses regarding the normal physiological function of Duffy, which include a potential role on red cells as a chemokine clearance factor to maintain chemokine gradients from tissue to blood (Horuk, 1994), and a role on endothelial cells in the presentation of chemokines to leukocytes (Hadley et al., 1994).

Presumably, fixation of the mutant Duffy allele in Africa was caused by selective pressure of *P. vivax* malaria. An opportunity to test this hypothesis prospectively may exist in Papua New Guinea, where *P. vivax* infection is endemic and where the identical promoter mutation has occurred independently on a different Duffy haplotype but is currently present at only a low frequency in the population (Zimmerman et al., 1999).

Chemokine and *Plasmodium* ligands have no significant sequence homology but cross-inhibit the binding of each other to Duffy. G α variants have been identified that block red cell invasion by *P. knowlesi* without activating CXCR1 or CXCR2 (Hesselgesser et al., 1995). A glutamic acid for alanine-substituted mutant named E6A is the most potent variant identified, but clinical development has not been reported. The Duffy-specific mAb Fy6 blocks chemokine binding (Horuk et al., 1993).

A murine ortholog of Duffy has been identified, and the gene is named *Dfy* (Luo et al., 1997; Tang et al., 1998). Features common with human Duffy include 63% aa identity; localization on mouse chromosome 1; expression in skeletal muscle, spleen, bone marrow, and brain; presence of an intron between codons 7 and 8; and a common chemokine binding profile. One difference is expression of the mouse gene in liver. Duffy-deficient mice have not yet been reported.

B. D6

The nomenclature for D6 has changed several times due to confusion about its signaling properties. Mouse D6 was originally identified as a bone marrow cDNA expressed in hematopoietic progenitor cells and placenta (Nibbs et al., 1997a). Ligands include multiple CC chemokines, including MIP-1 α , MCP-1, RANTES, and MCP-3, which were originally reported to induce calcium flux in transfected cells. Two human cDNAs were then reported that encode proteins 70% identical with mouse D6. Their sequences differ by 4 aa, which is thought to be due to allelic variation of the same gene. One variant was originally named "CCR9" in the GenBank database, but "D6" in the article describing it (Nibbs et al., 1997b). The other variant was named

"CCR10" (Bonini et al., 1997). Both variants were shown to bind similar CC chemokines as mouse D6, with the exception of MIP-1 α (see later), but signaling could not be demonstrated. Moreover, signaling for mouse D6 could not be reproduced (Nibbs et al., 1999). Therefore, the committee has not proposed a CCR designation for this molecule, and the groups originally responsible for its identification have agreed to use of D6 pending further research (Nibbs et al., 1999; J. Bonini, personal communication).

The difference in specificity of human and mouse D6 for MIP-1 α has been reconciled recently through the appreciation that there are two distinct but highly related MIP-1 α genes in human, named LD78 α and LD78 β , but only one in mouse. The LD78 α and LD78 β products have been tentatively named MIP-1 α S and MIP-1 α P, respectively, because of signature serine and proline residues that distinguish their structure and specificity for human D6 (Nibbs et al., 1999). In particular, MIP-1 α P is a selective agonist versus MIP-1 α S at human D6 in calcium flux assays. In vivo roles of D6 are undefined.

VIII. Virus-Encoded Chemokine Receptors

A. ECRF3

ECRF3 is another name for ORF 74 of Herpesvirus saimiri, a T lymphotropic γ -herpesvirus that is not pathogenic in its natural host, the squirrel monkey, but causes T cell transformation and acute leukemias and lymphomas in other primate hosts. The ECRF3 sequence was discovered as part of the saimiri genome sequencing project, and its relationship to chemokine receptors was identified through bioinformatics (Albrecht et al., 1992). When it is expressed in *Xenopus* oocytes, agonists include G α > IL-8 > NAP-2, which is the same specificity but a different hierarchy as for CXCR2, its closest known mammalian homolog (Ahuja and Murphy, 1993). Signaling includes calcium mobilization. Functional expression has not been successfully accomplished in mammalian cell lines, and there is no information about its biological role or expression in the context of virally infected cells.

B. US28

Human cytomegalovirus (HCMV) is a β -herpesvirus that causes life-threatening systemic infections in immunocompromised hosts, including patients with AIDS. UL33, US27, and US28 are ORFs in the unique long and unique short genomic regions of HCMV, respectively, that encode homologs of GPCRs (Chee et al., 1990). UL33 and US27 remain orphans, whereas US28 (30% aa identity to CCR1) has been demonstrated to bind the CC chemokines MIP-1 α , MIP-1 β , RANTES, MCP-1, and MCP-3 with similar affinity in transfected cells, as well as in the context of HCMV infection of human fibroblasts (Neote et al., 1993; Gao and Murphy, 1994; Bod-

aghi et al., 1998; Vieira et al., 1998). These chemokines are also US28 agonists, inducing calcium flux in both transfected and HCMV-infected cells, with RANTES having the highest potency (Gao et al., 1994; Vieira et al., 1998).

HCMV-infected fibroblasts are able to deplete endogenous RANTES and MCP-1 from media in a US28-dependent manner, suggesting a novel antichemokine mechanism for immune evasion (Bodaghi et al., 1998; Vieira et al., 1998). The CC chemokine specificity of US28 has recently been broadened to include fractalkine, which has suggested a possible adhesive role of US28, analogous to that of the cellular fractalkine receptor CX3CR1, that could mediate cell-cell spread of virus (Kledal et al., 1998). Consistent with this, US28 enhances cell-cell fusion by different viral proteins (Pleskoff et al., 1998). In this regard, US28 also has HIV coreceptor activity (Pleskoff et al., 1997), which has suggested a potential symbiotic relationship between the two viruses, with HIV providing immunosuppression needed for maximal HCMV replication and HCMV providing an additional mechanism for cell entry by HIV.

Recently, US28 was shown to transduce smooth muscle cell migration in response to RANTES and MCP-1 (Streblow et al., 1999). This provides a potential molecular mechanism to explain the link between smooth muscle cell infection by HCMV and vascular disease.

C. KSHV GPCR

KSHV GPCR is the product of ORF 74 of KS-associated herpesvirus or HHV8, a γ -herpesvirus and purported causal factor in KS, and two rare B cell neoplasms found in AIDS patients: primary effusion lymphoma and multicentric Castleman's disease (Cesarman and Knowles, 1999). KSHV GPCR is syntenic with ECRF3, the CXC chemokine receptor of Herpesvirus saimiri, and was discovered through bioinformatics as part of the HHV8 Genome Sequencing Project. It has been detected at the mRNA level in KS tissue and in B cell lymphomas (Cesarman and Knowles, 1999). Ligands include both CC and CXC chemokines, with highest apparent affinity for IL-8 (Arvanitakis et al., 1997). The receptor is unique among chemokine receptors in being constitutively active. Several chemokine ligands, most potently GRO α , can increase the activity in transfected mammalian cells and are therefore agonists (Gershengorn et al., 1998), whereas other chemokines, such as IP-10, vMIP-II, and SDF-1, decrease the activity and are inverse agonists (Geras-Raaka et al., 1998a,b; Rosenkilde et al., 1999). Signaling includes pertussis toxin-resistant phospholipase C activation in transfected COS cells and cell transformation. Activation of protein kinase C and cotransfection of GRK-5 have also caused down-regulation of activity (Geras-Raaka et al., 1998c). The transduction mechanism is not defined but does not appear to be G $_i$ as for other chemokine receptors. Biological function includes angiogenesis and oncogenesis

in transfected cells (Bais et al., 1998). KSHV GPCR may be responsible in part for KS pathogenesis. If so, the development of inverse agonists may be therapeutically useful.

D. UL12

The T lymphotropic β -herpesvirus HHV-6 causes acute and latent infections and encodes two GPCR homologs, U12 and U51. The U12 gene is expressed late in infection from a spliced mRNA, and when expressed in transfected cells, the encoded protein functions as a calcium-mobilizing CC chemokine receptor specific for RANTES, MIP-1 α , MIP-1 β , and MCP-1 (Isegawa et al., 1998). Function in the context of HHV6 infected cells and biological roles *in vivo* have not been delineated.

E. E1

ORF E1 of equine herpesvirus 2 has high sequence similarity to CC chemokine receptors (Telford et al., 1995) and was recently demonstrated to mediate calcium flux and chemotaxis in response to eotaxin (Carmada et al., 1999).

The subject of viral chemokine and orphan chemokine receptor-like proteins is quite large and interesting, but unfortunately is beyond the scope of this review, which is limited to 7TMD chemokine binding proteins and receptors. Information about putative viral chemokine receptors, viral chemokines, and non-7TMD viral chemokine binding proteins, some of which have already been demonstrated to be virulence factors, is summarized in Table 5 and in recent reviews (Pease and Murphy, 1998; Lalani and McFadden, 1999).

IX. Conclusions

The attention paid to chemokine receptors has greatly increased as AIDS has merged with inflammation in a common area of pharmacological opportunity. Nevertheless, notably absent from most receptor descriptions given here are lists of potent, selective antagonists typically found for other types of GPCRs and well-defined indications in disease. This is in part because the field is young and has had an inverted history relative to classic receptors, which began with antagonists and even approved drugs before the molecular basis of action was known. In the future this will no doubt change for chemokine receptors, but the molecular information reviewed here already indicates reason to anticipate significant, idiosyncratic hurdles, including major cross-species differences in both structure and repertoire of receptors and ligands; unanticipated difficulties in identifying lead compounds, which has already been encountered for CXCR1; and extensive redundancy in the system, with regard to both endogenous ligands, nonchemokine classic chemoattractant receptors (Murphy, 1994), and biology. As these complexities are sorted out, it is the sincere hope of the committee that the

nomenclature system described herein will help to prevent any ambiguity in communication about specific receptor targets.

Acknowledgments. We thank Josh Farber for critical reading of the manuscript and excellent suggestions.

REFERENCES

Agostini C, Casatella M, Zambello R, Trentin L, Gasperini S, Perin A, Piazza F, Siverio M, Faccio M, Dziejman M, Chilosi M, Qin S, Luster AD and Semenzato G (1998) Involvement of the IP-10 chemokine in sarcoid granulomatous reactions. *J Immunol* 161:6413–6420.

Ahuja SK, Ozolski T, Milatovich A, Francke U and Murphy PM (1992) Molecular evolution of the human interleukin-8 receptor gene cluster. *Nat Genet* 2:31–36.

Ahuja SK and Murphy PM (1993) Molecular polarity of mammalian interleukin-8 receptor type B for Hepervirus saimiri. *J Biol Chem* 268:20691–20694.

Ahuja SK, Lee JC and Murphy PM (1996) CXCR chemokines bind to unique sets of selectivity determinants that can function independently and are broadly distributed on multiple domains of human interleukin-8 receptor B: Determinants of high affinity binding and receptor activation are distinct. *J Biol Chem* 271:2255–2262.

Ahuja SK and Murphy PM (1996) The CXCR chemokines growth-regulated oncogene (CRO) alpha, GRO-beta, GRO-gamma, neutrophil-activating peptide-2, and epithelial cell-derived neutrophil-activating peptide-7 are potent agonists for the type B, but not the type A, human interleukin-8 receptor. *J Biol Chem* 271:20545–20550.

Altieri JC, Nicholas J, Biller D, Cameron KR, Biesinger B, Newman C, Wittmann S, Craxton MA, Coleman H, Fleckenstein B, et al. (1992) Primary structure of the Hepervirus saimiri genome. *J Virol* 66:5047–5058.

Althoff A, Ferriani S, Benelli R, Sforzini S, Giungio D, Aluigi MG, Proudfoot AE, Aluigi S, Wells TN, Mariani G, Rabin RL, Farber JM and Nozawa DM (1998) HIV-1 Tat protein mimics of chemokines. *Proc Natl Acad Sci USA* 95:11513–11518.

Alcami A, Symons JA, Collins PD, Williams TJ and Smith GL (1998) Blockade of chemokine activity by a soluble chemokine binding protein from vaccinia virus. *J Immunol* 160:2423–2433.

Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM and Berger EA (1996) CC CR2: A RANTES, MIP-1, MIP-1b receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science (Wash DC)* 272:1955–1958.

Amara A, Gull SL, Schwartz G, Mouton R, Mosier B, Loetscher P, Baggiolini M, Virelizier JL and Arenzana-Seisdedos F (1997) HIV coreceptor downregulation as antiviral principle: SDF-1alpha-dependent internalization of the chemokine receptor CXCR4 contributes to inhibition of HIV replication. *J Exp Med* 186:139–146.

Annunzio F, Cosmi L, Galli G, Beltrame C, Romagnani P, Manetti R, Romagnani S and Maggi E (1999) Assessment of chemokine receptor expression by human Th1 and Th2 cells in vitro and in vivo. *J Leukoc Biol* 65:691–699.

Ansel KM, McEvey-Williams LJ, Ngo VN, McEvey-Williams MG and Cyster JG (1999) In vivo-activated CD4 T cells upregulate CXCR chemokine receptor 5 and reprogram their response to lymphoid chemokines. *J Exp Med* 190:1123–1134.

Arenzana-Seisdedos F, Virelizier JL, Rousset D, Clark-Lewis I, Loetscher P, Moser B and Baggiolini M (1996) HIV blocked by chemokine antagonist. *Nature (Lond)* 383:400.

Arvanitaki L, Geras-Raska E, Varma A, Gershengorn MC and Cesarman E (1997) Human herpesvirus KSHV encodes a constitutively active G-protein-coupled receptor linked to cell proliferation. *Nature (Lond)* 385:347–350.

Baba M, Imai T, Nishimura M, Kakizaki M, Takagi S, Hieshima K, Nomiyama H and Yoshie O (1997) Identification of CXCR3, the specific receptor for a novel lymphocyte-directed CC chemokine, RANTES. *J Biol Chem* 272:14988–14998.

Baba M, Nishimura O, Kanazaki N, Okamoto M, Sawada H, Iizawa Y, Shiraishi M, Aramaki Y, Okonogi K, Ogawa Y, Meguro K and Fujino M (1999) A small-molecule, nonpeptide CXCR3 antagonist with highly potent and selective anti-HIV-1 activity. *Proc Natl Acad Sci USA* 96:7068–7070.

Baggiolini M, Dewald B and Moser B (1997) Human chemokines: An update. *Annu Rev Immunol* 15:675–705.

Baird JW, Nibbs RJ, Komai-Koma M, Connolly JA, Ottersbach K, Clark-Lewis I, Lew FY and Graham GJ (1999) ESKine, a novel β -chemokine, is differentially upregulated to produce secreted and nuclear targeted isoforms. *J Biol Chem* 274:33496–33503.

Bais C, Santomaso B, Coss O, Arvanitaki L, Raska EG, Gutkind JS, Asch AS, Cesarman E, Gershengorn MC and Mesri RA (1998) G-protein-coupled receptor of Kaposi's sarcoma-associated herpesvirus is a viral oncogene and angiogenesis activator. *Nature (Lond)* 391:58–59 [Published erratum appears in *Nature (Lond)* (1998) 392:210].

Baldwin JM (1993) The probable arrangement of the helices in G protein-coupled receptors. *EMBO J* 12:1683–1703.

Bareille L, Loetscher M, Tobler A, Baggiolini M and Moser B (1995) Sequence variation of a novel heptapeptide leukocyte receptor through alternative transcript formation. *Biochem J* 309:773–779.

Bazan HA, Alkhatib G, Broder CC and Berger EA (1998) Patterns of CXCR5, CXCR4, and CXCR3 usage by enveloped glycoproteins from human immunodeficiency virus type 1 primary isolates. *J Virol* 72:4855–4861.

Bazan JF, Bacon KB, Hardman G, Wang W, So K, Rossi D, Greaves DR, Zlotnik A and Schall TJ (1997) A new class of membrane bound chemokine with a CXCR3 motif. *Nature (Lond)* 385:64–64.

Beisser PS, Vink G, Van der Vliet AG, Guals G, Vanhoose SJ and Bruggeman CA (1998) The R33 G protein-coupled receptor gene of rat cytomegalovirus plays an essential role in the pathogenesis of viral infection. *J Virol* 72:2352–2363.

Ben-Baruch A, Xu L, Young PR, Bengali K, Oppenheim JD and Wang J-Z (1995)

Monocyte chemotactic protein-3 (MCP3) interacts with multiple leukocyte receptors: C-C CR1, a receptor for macrophage inflammatory protein-1 alpha/Rantes, is also a functional receptor for MCP3. *J Biol Chem* 270:22123–22128.

Benkiran M, Jin DY, Chua RF, Koup RA and Jeang KT (1997) Mechanism of transdominant inhibition of CXCR5-mediated HIV-1 infection by ccr5delta32. *J Biol Chem* 272:30603–30606.

Berger EA, Dong HW, Feng Y, Kober BTM, Littman DR, Moore JW, Sautentat GJ, Schuitemaker H, Sodroski J and Weiss RA (1998) A new classification for HIV-1. *Nature (Lond)* 391:240.

Berger EA, Murphy PM and Farber JM (1999) Chemokine receptors as HIV-1 coreceptors: Roles in viral entry, tropism and disease. *Annu Rev Immunol* 17:507–709.

Bernardini G, Hedrick J, Sozzani S, Luitl W, Spinetti G, Weiss M, Menon S, Zlotnik A, Mantovani A, Santoni A and Napolitano M (1998) Identification of the CC chemokines TARC and macrophage inflammatory protein-1 beta as novel functional ligands for the CXCR receptor. *Eur J Immunol* 28:5858–5868.

Birkbach M, Joseph K, Yamashita H, Lenoir G and Kiehl E (1993) Epstein-Barr virus-induced genes: First lymphocyte-specific G-protein-coupled peptide receptors. *J Virol* 67:2209–2220.

Blanchard C, Migonette I, Lee B, Vakili J, Doranz BJ, Govaerts C, Vassart G, Doms RW and Parmentier M (1999) CXCR5 binds multiple CC-chemokines: MCP-3 acts as a natural antagonist. *Blood* 94:1899–1905.

Bleul CC, Farzan M, Choe H, Parolin C, Clark-Lewis I, Sodroski J and Springer TA (1996) The lymphocyte chemottractant SDF-1 is a ligand for LEST/FRUS and blocks HIV-1 entry. *Nature (Lond)* 382:830–833.

Bleul CC, Wu L, Henry M, Clark-Lewis I, Springer TA and Mackay CR (1997) The HIV coreceptors CXCR4 and CXCR5 are differentially expressed and regulated on human T lymphocytes. *Proc Natl Acad Sci USA* 94:1926–1930.

Bodaghi B, Jones TR, Zipeto D, Vita C, Sun L, Laurent L, Arenzana-Seisdedos F, Virelizier JL and Michelon S (1998) Chemokine sequestration by viral chemokine receptors as a novel viral escape strategy: withdrawal of chemokines from the environment of cytomegalovirus-infected cells. *J Exp Med* 188:855–866.

Boisvert WA, Santiago R, Curtiss LK and Terkeltaub RA (1998) A leukocyte homologue of the IL-8 receptor CXCR2 mediates the accumulation of macrophages in atherosclerotic lesions of LDL receptor-deficient mice. *J Clin Invest* 101:935–943.

Bonacchi R, Bianchi G, Bordignon PP, D'Ambrosio D, Lang R, Borzatti A, Sozzani S, Allavena P, Gray PA, Mantovani A and Sinigaglia F (1998) Differential expression of chemokine receptors and chemotactic responsiveness of type 1 T helper cells (Th1) and Th2. *J Exp Med* 187:189–198.

Bonini JA, Martin SK, Dralyuk P, Roe MW, Philipson LH and Steiner DF (1997) Cloning, expression, and chromosomal mapping of a novel human CC-chemokine receptor (CXCR10) that displays high-affinity binding for MCP-1 and MCP-3. *DNA Cell Biol* 17:101–110.

Boring L, Gossling J, Chenese SW, Kunkel SL, Farsse RV Jr, Broxmeyer HE and Charo IF (1997) Impaired monocyte migration and reduced type 1 (Th1) cytokine responses in C-C chemokine receptor 2 knockout mice. *J Clin Invest* 100:2552–2561.

Boring L, Gossling J, Clerly M and Charo IF (1998) Decreased lesion formation in CXCR2-/- mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature (Lond)* 394:894–897.

Boring L, Gossling J, Montecarlo FS, Lusis AJ, Teou C-L and Charo IF (1996) Molecular cloning and functional expression of murine IL-8 (monocyte chemoattractant protein 1) and murine macrophage inflammatory protein 1 receptors: Evidence for two closely linked C-C chemokine receptors on chromosome 9. *J Biol Chem* 271:7551–7558.

Bouffard F, Endo Y, Collins PD, Takeuchi Y, Reeves JD, Schweickart VL, Siani MA, Sasaki T, Williams TJ, Gray PW, Moore PS, Chang Y and Weiss RA (1997) Angiogenic and HIV-inhibitory functions of KSHV-encoded chemokines. *Science* 278:290–294.

Bozic CR, Gerard NP, von Uexkull-Guldenband C, Kolakowski LF Jr, Conklyn MJ, Berman J, Croucher BJ and Gerard C (1994) The murine interleukin-8 type B receptor homologue and its ligands. *J Biol Chem* 269:29355–29358.

Bridger GB, Skerj R, Padmanabhan S, Martellucci SA, Henson GW, Struyf S, Witvrouw M, Scholz D and De Clercq E (1999) Synthesis and structure-activity relationships of phenylacetylenyl(methylene)-linked bis-azamacrocycles that inhibit HIV-1 and HIV-2 replication by antagonism of the chemokine receptor CXCR4. *J Med Chem* 42:3917–3931.

Broadbent VC, Boylan AM, Hoeffel JM, Kim KJ, Sadick M, Chuntharapal A and Robert CA (1994) Neutralization of IL-8 inhibits neutrophil influx in a rabbit model of endotoxin-induced pleurisy. *J Immunol* 152:2949–2957.

Broxmeyer HE, Cooper S, Caccalano G, Hogue NL, Balish E and Moore MW (1996) Involvement of Interleukin (IL) 8 receptor in negative regulation of myeloid progenitor cells in vivo: Evidence from mice lacking the murine IL-8 receptor homolog. *J Exp Med* 184:1829–1839.

Broxmeyer HE, Cooper S, Hange G, Gao JL and Murphy PM (1999) Dominant myelopoietic effector functions mediated by chemokine receptor CXCR1. *J Exp Med* 190:1987–1992.

Broxmeyer HE, Mantel CR and Aronica SM (1997) Biology and mechanisms of action of xenobiotic stimuli and myeloid progenitor cell proliferation and suppression by chemokines. *Stem Cells* 15 (suppl 1):69–77.

Burgstahler R, Kempeks B, Steube K and Lipp M (1995) Expression of the chemokine receptor BLR2/EBH1 is specifically transactivated by Epstein-Barr virus. *Proc Natl Acad Sci USA* 92:1317–1321.

Caccalano G, Lee J, Kikly K, Rhee AM, Pitta-Meck S, Hultgren B, Wood WJ and Moore MW (1994) Regulation of B cell expansion in mice that lack the murine IL-8 receptor homolog. *Science (Wash DC)* 265:682–685.

Camarda G, Spinetti G, Bernardini G, Malc R, Davis-Poynter N, Capogrossi MC and Napolitano M (1999) The human herpesvirus 2 EI open reading frame encodes a functional chemokine receptor. *J Virol* 73:9843–9848.

Campbell EM, Charo IF, Kunkel SL, Strieter RM, Boring L, Gossling J and Lukacs NW (1999) Monocyte chemoattractant protein-1 mediates cockroach allergen-

- induced bronchial hyperactivity in normal but not $\text{CCR2}^{-/-}$ mice: The role of mast cells. *J Immunol* 163:2160-2167.
- Campbell JH, Bowman EP, Murphy K, Youngman KR, Siani MA, Thompson DA, Wu L, Zlotnik A, and Butcher EC (1998) 6-C-kine (SLC), a lymphocyte adhesion-triggering chemokine expressed by high endothelium, is an agonist for the MIP-3beta receptor CCR7 . *J Cell Biol* 141:1053-1059.
- Campbell JH, Haraldson G, Pan J, Rotstein J, Qin S, Ponath P, Andrew DP, Warnke R, Ruffing N, Kassam N, Wu L, and Butcher EC (1998) Chemokine receptor CCR4 in vascular recirculation by cutaneous but not intestinal memory T cells. *Nature (Lond)* 400:767-770.
- Campbell JH, Hadravsky J, Zlotnik A, Siani MA, Thompson DA and Butcher EC (1998) Chemokines and the arrest of lymphocytes rolling under flow conditions. *Science* 279:381-384.
- Caro JX, Gershon PD and Black DN (1995) Sequence analysis of *HindIII* Q2 fragment of capripoxvirus reveals a putative gene encoding a G-protein-coupled chemokine receptor homologue. *Virology* 206:207-212.
- Cesmaner E and Knowles DM (1999) The role of Kaposi's sarcoma-associated herpesvirus (KSHV/HHV-8) in lymphoproliferative diseases. *Semin Cancer Biol* 9:165-174.
- Charbonnier AS, Kohrgruber N, Kriebelner E, Stügel G, Rot A and Mourer D (1999) Macrophage inflammatory protein 3a is involved in the constitutive trafficking of epidermal Langerhans cells. *J Exp Med* 191:1753-1768.
- Churo IF, Myers SJ, Herman A, Franci C, Connolly AJ and Coughlin SR (1994) Molecular cloning and functional expression of two monocyte chemoattractant protein 1 receptors reveals alternative splicing of the carboxyl-terminal tails. *Proc Natl Acad Sci USA* 91:2752-2756.
- Chaudhuri A, Nielsen S, Eklund J, Zbrenska V, Fang F and Pogo AO (1997) Detection of Duffy antigen in the plasma membranes and caveolae of vascular endothelial and epithelial cells of nonerythroid origins. *Blood* 89:701-712.
- Chaudhuri A, Polyakova J, Zbrenska V and Pogo AO (1995) The coding sequence of Duffy blood group gene in humans and simians: Restriction fragment length polymorphism, antibody and malpian parasitic specificity, and expression in nonerythroid tissues in Duffy-negative individuals. *Blood* 85:615-621.
- Chaudhuri A, Polyakova J, Zbrenska V, Williams K, Galatius X and Pogo AO (1993) Cloning of glycoprotein D cDNA, which encodes the major subunit of the Duffy blood group system and the receptor for the Plasmodium vivax malaria parasite. *Proc Natl Acad Sci USA* 90:10795-10797.
- Chaudhuri A, Zbrenska V, Polyakova J, Pogo AO, Hesselgesell J and Horuk R (1994) Expression of the Duffy antigen in K562 cells: Evidence that it is the human erythrocyte chemokine receptor. *J Biol Chem* 269:7835-7838.
- Chee MS, Satchwell SC, Frederix B, Wente KM and Soderstrom JG (1990) Human erythrocyte antigen encodes three G protein-coupled receptor homologues. *Nature (Lond)* 344:774-777.
- Chen JD, Bai X, Yang AG, Cong Y and Chen SY (1997) Inactivation of HIV-1 chemokine co-receptor CXCR-4 by a novel intraline strategy. *Nat Med* 3:1110-1116.
- Chitnis CE and Miller LH (1994) Identification of the erythrocyte binding domains of Plasmodium vivax and Plasmodium knowlesi proteins involved in erythrocyte invasion. *J Exp Med* 180:497-506.
- Choe H, Farzan M, Konkel M, Martin K, Sun Y, Marcon L, Cayabyab M, Berman M, Dorf ME, Gerard N, Gerard C and Sodroski J (1998) The orphan seven-transmembrane receptor A9 supports the entry of primary T-cell-line-tropic and dual-tropic human immunodeficiency virus type 1. *J Virol* 72:6113-6118.
- Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, Wu L, Mackay CR, LaRosa G, Newman W, Gerard NP, Gerard C and Sodroski J (1996) The β -chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV isolates. *Cell* 85:1135-1148.
- Chuntharapai A, Lee J, Hebert CA and Kim KJ (1994) Monoclonal antibodies detect different distribution patterns of IL-8 receptor A and IL-8 receptor B on human peripheral blood leukocytes. *J Immunol* 153:662-668.
- Clark-Lewis I, Kim KS, Rajarathnam K, Gong JH, Dewald B, Moser B, Baggiolini M and Smyth BD (1995) Structure-activity relationships of chemokines. *J Leukoc Biol* 57:703-711.
- Clemens MJ, Marshall E, Durig J, Watanabe K, Howell A, Miles D, Earl H, Kiernan J, Griffiths A, Towson K, DeTaktis P, Teata NG, Dougal M, Hunter MG, Wood LM, Czaplewski LG, Millar A, Dexter TM and Lord BI (1998) A randomized phase-II study of BB-1001 (macrophage inflammatory protein-1alpha) in patients with advanced breast cancer receiving 5-fluorouracil, adriamycin, and cyclophosphamide chemotherapy. *Breast* 9:2132-2140.
- Clore GM and Gronenborn AM (1995) Three-dimensional structures of alpha and beta chemokines. *FASEB J* 9:57-62.
- Cocchi F, DeVito AL, Garzino-Demo A, Arya SK, Gallo RC and Lusso P (1995) Identification of RANTES, MIP-1a, and MIP-1b as major HIV suppressive factors produced by T-tropic macrophages. *Science* 267:1631-1635.
- Cole KE, Strick CA, Paradis TJ, Opbrone KT, Loetscher M, Gluders RP, Lin W, Boyd JG, Moser B, Wood DE, Sahagan BG and Neote K (1998) Interferon-inducible T cell alpha chemokine (ITAC) a novel non-ELR CC chemokine with potent activity on activated T cells through selective high affinity binding to CXCR3. *J Exp Med* 187:2019-2027.
- Combadere C, Ahuja SK and Murphy PM (1995a) Cloning and functional expression of a human eosinophil CC chemokine receptor. *J Biol Chem* 270:16491-16494 [erratum published in *J Biol Chem* (1995) 270:30263].
- Combadere C, Ahuja SK and Murphy PM (1995b) Cloning, chromosomal localization, and RNA expression of a human beta chemokine receptor-like gene. *DNA Cell Biol* 14:673-680.
- Combadere C, Ahuja SK, Tiffany HL and Murphy PM (1996) Cloning and functional expression of CC CKR3, a human monocyte CC chemokine receptor selective for MIP-1a, MIP-1b and RANTES. *J Biol Chem* 271:1517-1521.
- Combadere C, Gao J, Tiffany HL and Murphy PM (1998a) Gene cloning, RNA distribution, and functional expression of mCXCR1, a mouse chemotactic receptor for the CX3C chemokine fractalkine. *Biochem Biophys Res Commun* 253:728-732.
- Combadere C, Salzwedel K, Smith ED, Tiffany HL, Berger EA and Murphy PM (1998b) Identification of CX3CR1: A chemotactic receptor for the human CX3C chemokine fractalkine, and a fusion coreceptor for HIV-1. *J Biol Chem* 273:23799-23804.
- Cook DN, Beck MA, Coffman TM, Kirby SI, Sheridan JF, Pragnell IB and Smithies O (1998) Requirement of MIP-1a for an inflammatory response to viral infection. *Science (Wash DC)* 280:1583-1585.
- Coulfin F, Power CA, Alouani S, Peitsch MC, Schroeder JM, Moshizuki M, Clark-Lewis I and Wells TM (1997) Characterisation of macrophage inflammatory protein-1 (MIP-1) as a member of the macrophage-inflammatory-protein family of chemokines. *Eur J Biochem* 248:7-15.
- Crump MP, Gong JH, Loetscher P, Rajarathnam K, Amara A, Arenzana-Seisdedos F, Virelizier JM, Baggiolini M, Sykes BD and Clark-Lewis I (1997) Solution structure and basis for functional activity of stromal cell-derived factor-1: dissociation of CXCR4 activation from binding and inhibition of HIV-1. *EMBO J* 16:6996-7007.
- Cyster JG (1999a) Chemokines and the homing of dendritic cells to the T cell areas of lymphoid organs. *J Exp Med* 189:447-450.
- Cyster JG (1999b) Chemokines and cell migration in secondary lymphoid organs. *Science* 286:2082-2086.
- Daelmolen D, Schols D, Witvrouw M, Pannecouque C, Hase S, Van Dooren S, Hamy F, Kikmakai T, de Clercq E and VanDamme AM (2000) A second target for the peptidate Tat/transportation response element inhibitor CPG4222: Inhibition of human immunodeficiency virus replication by a CXCR4-chemokine receptor 4-mediated virus entry. *Mol Pharmacol* 57:116-124.
- Dairaghi DJ, Fan RA, McMaster BE, Hanley MR and Schall TJ (1999) HIV-1-induced VAMP1 selectively engages chemokine receptor CXCR4: Agonist and antagonist profiles of viral chemokines. *J Biol Chem* 274:21569-21574.
- Dairaghi DJ, Oldham OB, Bacon KB and Schall TJ (1997) Chemokine receptor CXCR3 function is highly dependent on local pH and ionic strength. *J Biol Chem* 272:28206-28209.
- Damon I, Murphy PM and Moss B (1998) Broad spectrum chemokine antagonistic activity of a human poxvirus chemokine homologue. *Proc Natl Acad Sci USA* 95: 6403-6407.
- Daniels GD, Zou J, Charlemagne J, Partula S, Cunningham C and Seemayer CE (1999) Cloning of two chemokine receptor homologs (CXCR-4 and CXCR-7) in zebrafish *Oreochromis mossambicus*. *J Leukoc Biol* 65:684-690.
- Darmonne WC, Rice GC, Mohler MA, Apple T, Hebert CA, Valente AJ and Baker JB (1991) Red blood cells are a sink for interleukin 8, a leukocyte chemotaxin. *J Clin Invest* 88:1362-1369.
- Daugherly BL, Siciliano SJ, DeMartino JA, Malkowski L, Sirodina A and Springer MS (1996) Cloning, expression, and characterization of the human eosinophil eotaxin receptor. *J Exp Med* 183:2349-2354.
- Davis-Poynter NJ, Lynch DM, Vally H, Shellam GR, Rawlinson WD, Barrell BG and Farrell HE (1997) Identification and characterization of a G protein-coupled receptor for human eotaxin, a chemotactic factor for murine eosinophils. *J Virol* 71:1511-1520.
- Dean M, Carrington M, Winkler C, Hutley GA, Smith WM, Allikmets R, Goeder J, Buchbinder SP, Vittinghoff E, Gomperts K, Donfield S, Vlahov D, Kaslow R, Sah A, Rinaldo C, Detels R, Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE and O'Brien SJ (1996) Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CXCR5 structural gene. *Science (Wash DC)* 273:1856-1862.
- Deng H, Liu R, Rillemer W, Choe S, Uzunutaz D, Burkhardt M, Dimarzio P, Marnett S, Sutton RE, Hill CM, Littman D and Littman DR (1996) Identification of a major co-receptor for primary isolates of HIV-1. *Nature (Lond)* 381:661-666.
- Deng HK, Uzunutaz D, Kewal Ramani VN and Littman DR (1997) Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature (Lond)* 388:296-300.
- Dean MC, Vanvubert B, Vicari A, Bridon JM, Oldham E, Ais-Yahia S, Briere F, Zlotnik A, Lebeque S and Caux C (1998) Selective recruitment of immature and mature dendritic cells by distinct chemokines expressed in different anatomic sites. *J Exp Med* 186:373-386.
- Dell'Acqua D, Grossi F, Farber K, Sharon MP, Mei SH, Gost JB, Daar ES, Doms RW and O'Brien WA (1997) A small-molecule inhibitor directed against the chemokine receptor CXCR4 prevents its use as an HIV-1 coreceptor. *J Exp Med* 186:1395-1400.
- Dobner T, Wolf I, Ehrlich T and Lipp M (1992) Differentiation-specific expression of a novel β -chemokine coreceptor for Burkitt's lymphoma. *Eur J Immunol* 22:2785-2789.
- Donzella GA, Schols D, Lin SW, Este JA, Nagashima KA, Madden PJ, Allaway GP, Sukumar TP, Henson G, De Clercq E and Moore JP (1998) AMD3100, a small molecule inhibitor of HIV-1 entry via the CXCR4 coreceptor. *Nat Med* 4:72-77.
- Dorner B, Gao J, Sharrow KA, Sharon MP, Mei SH, Gost JB, Daar ES, Doms RW and O'Brien WA (1997) A small-molecule inhibitor directed against the chemokine receptor CXCR4 prevents its use as an HIV-1 coreceptor. *J Exp Med* 186:1395-1400.
- Dorand B, Rucker J, Yi Y, Smyth RJ, Sansom M, Peiper SC, Parmentier M, Collman RC and Doms RW (1994) A dual tropic primary HIV-1 isolate that uses fusin and the β -chemokine receptors CXCR-5, CXCR-3, and CXCR-2b as fusion co-factors. *Cell* 86:1149-1158.
- Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, Canavan C, Broder CC, Broder JP, Paxton WA, Moore JP and Paxton WA (1996) HIV-1 entry into CD4 cells is mediated by the chemokine receptor CXCR-5. *Nature (Lond)* 381:667-673.
- Dunstan AN, Salfanra MN, Adhikari S, Xia Y, Feng L and Harrison JK (1996) Identification of two genes orthologous to the human interleukin-8 receptors. *Biol. Chem. Biophys.* 271:32770-32778.
- Endres MJ, Clapham PR, Marsh M, Ahuja M, Turner JD, McKnight A, Thomas JF, Stoebenau-Haggarty B, Choe S, Vance PD, Wells TN, Power CA, Suterवास SS,

- Doms RW, Leland NR and Hoxie JA (1996) CD4-independent infection by HIV-2 is mediated by fusin/CXCR4. *Cell* 87:745–756.
- Endres MJ, Gurley CA, Gao XN, Li L and Hedrick JA (1999) The Kaposi's sarcoma-associated herpesvirus (KSHV)-encoded chemokine vMIP-1 is a specific agonist for the CC chemokine receptor (CCR8). *J Exp Med* 189:1993–1998.
- Farzani M, Choe H, Martin K, Maroon L, Hofmann W, Karlsson C, Sun Y, Barrett P, Marchand N, Sullivan N, Gerard N, Gerard C and Soderstrom J (1997) Two orphan seven-transmembrane segment receptors which are expressed in CD4-positive cells support simian immunodeficiency virus infection. *J Exp Med* 186:405–411.
- Farzani M, Mirzabekov T, Kolchinsky P, Wyatt R, Cayabach M, Gerard NP, Gerard C, Soderstrom J and Choe H (1999) Tyrosine sulfation of the amino terminus of CCR8 facilitates HIV-1 entry. *Cell* 85:659–670.
- Feng L, Chen S, Garcia GE, Xia Y, Sliwa MA, Boti P, Wilson CR, Harrison JC and Bacon KB (1999) Prevention of crescentic glomerulonephritis by immunoneutralization of the fractalkine receptor CX3CR1. *Kidney Int* 56:612–620.
- Feng Y, Broder CC, Kennedy PE and Berger E (1996) HIV-1 entry co-factor: functional cDNA cloning of a seven-transmembrane G-protein coupled receptor. *Science* (Wash DC) 272:872–877.
- Fleming P, Davis-Poynter N, Degli-Esposti M, Densley E, Papadimitriou J, Shillam G and Farrell H (1999) The murine cytomegalovirus chemokine homolog, m121, is a determinant of viral pathogenicity. *J Virol* 73:6800–6809.
- Fong AM, Robinson LA, Stecher DA, Toddler TF, Yoshie O, Imai T and Patel DD (1998) Fractalkine and CX3CR1 mediate a novel mechanism of leukocyte capture, firm adhesion, and activation under physiologic flow. *J Exp Med* 188:1413–1419.
- Forsmann U, Ugucioni M, Loetscher P, Dahinden CA, Langen H, Thelen M and Baggiolini M (1997) RANTES-2, a novel CC chemokine that is selective for the chemokine receptor CXCR5, and like RANTES on human eosinophil and basophil leukocytes. *J Exp Med* 185:2171–2176.
- Forster R, Enrich T, Kremmer E and Lipp M (1994) Expression of the G-protein-coupled receptor BLR1 defines mature, recirculating B cells and a subset of T-helper memory cells. *Immunity* 8:839–849.
- Forster R, Mattis AE, Kremmer E, Wolf E, Brem G and Lipp M (1996) A putative chemokine receptor, BLR1, directs B cell migration to defined lymphoid organs and specific anatomic compartments of the spleen. *Cell* 87:1037–1047.
- Forster R, Schubel A, Breitfeld D, Kremmer R, Renner-Muller I, Wolf E and Lipp M (1999) CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* 99:23–33.
- Foxman EF, Campbell JJ and Butcher EC (1997) Multistep navigation and the combinatorial control of leukocyte chemotaxis. *J Cell Biol* 139:349–360.
- Frade JM, Mellado M, del Real G, Gutierrez-Ibanez JC, Lind P and Martinez-A C (1997) Characterization of the CCR2 chemokine receptor: Functional CCR2 receptor expression in B cells. *J Immunol* 159:5576–5584.
- Frodi R, Gierachik P and Moegge B (1998) Genomic organization and expression of the CXCR4 gene in mouse and man: Absence of a splice variant corresponding to mouse CXCR4-B in human tissue. *J Recept and Transmembr* 18:321–344.
- Gallo RC (1999) Tat as one key to HIV-induced immune pathogenesis and Tat (correction of Pat) toxigen as an important component of a vaccine. *Proc Natl Acad Sci USA* 96:8324–8326.
- Gao JL, Kuhns DS, Tiffany HL, McInnes D, Li X, Franke U and Murphy PM (1998) Structure and functional expression of the human macrophage inflammatory protein 1 alpha/RANTES receptor. *J Exp Med* 177:1421–1427.
- Gao JL and Murphy PM (1994) Human cytomegalovirus open reading frame US28 encodes a functional beta chemokine receptor. *J Biol Chem* 269:28539–28542.
- Gao JL and Murphy PM (1995) Cloning and differential tissue-specific expression of three mouse β chemokine receptor-like genes, including the gene for a functional MIP-1 receptor. *J Biol Chem* 270:17494–17501.
- Gao JL, Sen AL, Kitaura M, Yoshie O, Rothenberg ME, Murphy PM and Luster AD (1996) Identification of a mouse eosinophil receptor for the CC chemokine eotaxin. *Biochem Biophys Res Commun* 223:679–684.
- Gao JL, Wynn TA, Chang Y, Lee EJ, Broxmeyer HE, Cooper S, Tiffany HL, Westphal H, Kwon-Chung J and Murphy PM (1997) Impaired host defense, hematopoiesis, granulomatous inflammation and type 1-type 2 cytokine balance in mice lacking CC chemokine receptor 1. *J Exp Med* 186:1595–1606.
- Gao W, Topfman PS, King AJ, Smiley ST, Cismasini V, Lu B, Gerard CJ and Hancock WW (2000) Targeting of the chemokine receptor CCR1 suppresses development of acute and chronic cardiac allograft rejection. *J Clin Invest* 105:335–44.
- Garcia-Zepeda EA, Combadiere C, Rothenberg ME, Srafi NN, Lavigne F, Hamid Q, Murphy PM and Luster AD (1998) Human monocyte chemoattractant protein-1 (MCP-1): A novel CC chemokine with activities on monocytes, eosinophils, and basophils induced in allergic and non-allergic inflammation that signals through the CC chemokine receptors CCR2-B and CCR3-B. *J Immunol* 157:5613–5626.
- Gerard C (1999) Understanding chemokine biology through mouse models: Riddles and answers. In *Chemokines in Disease* (Herbert CA ed) pp 44–52, Humana, Totowa, NJ.
- Gerard C, Frossard JL, Bhatia M, Saluja A, Gerard NP, Lu B and Steer M (1997) Targeted disruption of the beta-chemokine receptor CCR1 protects against pancreatitis-associated lung injury. *J Clin Invest* 100:222–231.
- Geras-Raska E, Varona A, Clark-Lewis I and Gershengorn MC (1998a) Kaposi's sarcoma-associated herpesvirus (KSHV) chemokine vMIP-2 and human SDF-1 α induce signaling by KSHV G-protein-coupled receptor. *Biochem Biophys Res Commun* 253:729–737.
- Geras-Raska E, Varona A, Clark-Lewis I and Gershengorn MC (1998b) Human interferon gamma-inducible protein 10 (IP-10) inhibits constitutive signaling of Kaposi's sarcoma-associated herpesvirus G-protein-coupled receptor. *J Exp Med* 188:405–408.
- Geras-Raska E, Arvanitaki I, Bais C, Cesarman E, Meiri EA and Gershengorn MC (1998c) Inhibition of constitutive SDF-1 α receptor signaling by a G-protein-coupled receptor by protein kinase C in mammalian cells in culture. *J Exp Med* 187:801–806.
- Gerber BO, Zanni MP, Ugucioni M, Loetscher M, Mackay CR, Pichler WJ, Yawalkar N, Baggiolini M and Moser B (1997) Functional expression of the receptor for CCR3 in T lymphocytes co-localizing with eosinophils. *Curr Biol* 7:836–843.
- Gershengorn MC, Varona A, Yarns A and Clark-Lewis I (1998) Chemokines activate Kaposi's sarcoma-associated herpesvirus G-protein-coupled receptor in mammalian cells in culture. *J Clin Invest* 102:1469–1472.
- Gerszten RE, Garcia-Zepeda EA, Lim YC, Yoshida M, Ding HA, Gimbrone MA Jr, Luster AD, Luscinskas FW and Rosenzweig A (1995) MCP-1 and IL-8 trigger firm adhesion of monocytes to vascular endothelium under flow conditions. *Nature* 378:118–123.
- Golla R and Banerjee AC (1998) Sequence specific cleavage of the HIV-1 coreceptor CCR5 gene by a hammer-head ribozyme and a DNA-enzyme: Inhibition of the coreceptor function by a DNA-enzyme. *FEBS Lett* 436:253–257.
- Gong W, Howard OM, Turpin JA, Grimm MC, Ueda H, Gray PW, Raport JC, Oppenheim JJ and Wang JM (1998) Monocyte chemotactic protein-2 activates CCR5 and blocks CXCR5-mediated HIV-1 entry/replication. *J Biol Chem* 273:4289–4292.
- Gong X, Gong W, Kuhns DS, Ben-Baruch A, Howard OM and Wang JM (1997) Monocyte chemotactic protein-2 (MCP-2) uses CCR1 and CCR2B as its functional receptors. *J Biol Chem* 272:11682–11685.
- Gonzalez JA, Pan Y, Lloyd CM, Jia GQ, Yu G, Dussault JB, Powers CA, Proudfoot AE, Coyle AJ, Gearing D and Gutierrez-Ramos JC (1999) Mouse monocyte-derived chemokine is involved in airway hyperreactivity and lung inflammation. *J Immunol* 163:403–411.
- Gosling J, Slaymaker S, Gu L, Tseng S, Zlot CH, Young SG, Rollins BJ and Charo IF (1999) MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B. *J Clin Invest* 103:733–741.
- Goya I, Gutierrez J, Varona R, Kremer L, Zaballos A and Marquez G (1998) Identification of CCR8 as the specific receptor for the human beta-chemokine 1309: cloning and molecular characterization of murine CCR8 as the receptor for TCA-3. *J Immunol* 160:1919–1981.
- Graham KA, Lalani AS, Macen J, Ness TL, Barry M, Liu LY, Lucas A, Clark-Lewis I, Moyer RW and McFadden G (1997) The T155kDa family of poxvirus-secreted proteins bind chemokines and modulate leukocyte influx into virus-infected tissues. *Virology* 222:13–24.
- Grossi D, Piper R, Pape M, Chen D, Wei Y, Iadelli F, O'Doherty U, Paxton W, Koup R, Mojsov S, Bhargava N, Clark-Lewis I, Baggiolini M and Steinman RM (1996) Efficient interaction of HIV-1 with purified dendritic cells via multiple chemokine coreceptors. *J Exp Med* 184:2433–2438.
- Groves DF, Wang W, Dairaghi DJ, Dine MC, Salas-Via B, Franz-Bacon K, Rossi D, Caux C, McClanahan T, Gordon S, Zlotnik A and Schall TJ (1997) CCR6, a chemokine receptor that interacts with macrophage inflammatory protein 3alpha and is highly expressed in human dendritic cells. *J Exp Med* 186:837–844.
- Green SP, Chakrabarti A and Curran TJ (1998) Interleukin-8 (IL-8) and stromal cell derived factor-1 (SDF-1) are chemokines that activate distinct signaling pathways to stimulate priming of the neutrophil NADPH oxidase through the type A or type B IL-8 receptor. *J Biol Chem* 273:25400–25405.
- Grimaldi JC, Yu XN, Grunig G, Seymour BW, Cottrez F, Robinson DS, Hosken N, Ferlin W, Wu Y, Soto H, O'Garra A, Howard MC and Coffman RL (1999) Depletion of eosinophils in mice through the use of antibodies specific for C-C chemokine receptor 3 (CCR3). *J Leuk Biol* 65:846–853.
- Gu L, Okada Y, Clinton SK, Gerard C, Sukhova GK, Libby P and Rollins BJ (1998) Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor-deficient mice. *Mol Cell* 2:725–731.
- Gunn MD, Kywau S, Tam C, Kakiuchi T, Matsuzawa A, Williams LT and Nakano H (1999) Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J Exp Med* 189:451–460.
- Gunn MD, Ngo VN, Ansel KM, Eklund EH, Cyster JG and Williams LT (1998a) A B-cell-homing chemokine made in lymphoid follicles activates Burkitt's lymphoma receptor-1. *Nature (Lond)* 391:799–803.
- Gunn MD, Kywau S, Tam C, Cyster JG, Rosen SD and Williams LT (1998b) A chemokine expressed in lymphoid high endothelial venules promotes the adhesion and chemotaxis of naive T lymphocytes. *Proc Natl Acad Sci USA* 95:258–263.
- Gupta SK, Lyvko PG, Pillarsetti K, Ohlstein E and Stadel JM (1998b) Chemokine receptors in human endothelial cells: Functional expression of CXCR4 and its transcriptional regulation by inflammatory cytokines. *J Biol Chem* 273:4282–4287.
- Gupta SK and Pillarsetti K (1999) Cutting edge: CXCR4-LO: Molecular cloning and functional expression of a novel human CXCR4 splice variant. *J Immunol* 163:2288–2292.
- Gupta SK, Pillarsetti K, Gray SL and Stadel JM (1998a) Molecular cloning of a novel chemokine receptor-like gene from early stage chick embryos. *Biochem Mol Int* 44:673–681.
- Haidis F, Viorard V, Autran B, Clark-Lewis I, Baggiolini M and Debs P (1998) HIV-specific T cell cytotoxicity mediated by RANTES via the chemokine receptor CCR3. *J Exp Med* 188:609–614.
- Hadley TJ, Lu ZH, Wasniowska K, Martin AW, Peliper SC, Hesselgeser J and Horuk R (1994) Postreceptor tyrosine endothelial cells in kidney express a multiplexed chemokine receptor that is structurally and functionally identical to the erythroid isoform, which is the Duffy blood group antigen. *J Clin Invest* 94:955–961.
- Hamada T, Mohle R, Hesselgeser J, Hoxie J, Nachman RL, Moore MA and Rafii S (1998) Transendothelial migration of megakaryocytes in response to stromal cell derived factor-1 (SDF-1) by human endothelial cells. *J Exp Med* 188:539–548.
- Hammond MR, Lapointe GR, Fackler HJ, Hibi T, Gallages CC, Gordon CA, Giedlin MA, Mullenbach G and Tekamp-Olson P (1995) IL-8 induces neutrophil chemotaxis predominantly via type 1 IL-8 receptors. *J Immunol* 155:1428–1433.
- Harrison JK, Barber CM and Lynch KR (1994) cDNA cloning of a G-protein-coupled receptor with a spinal cord and brain related to chemokine receptors. *Neurosci Lett* 169:85–89.
- Harrison JK, Jiang Y, Chen S, Xia Y, Maciejewski D, McNamara RK, Streit WJ, Salafrafrance MN, Adhikari S, Thompson DA, Boti P, Bacon KB and Feng L (1998)

- Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proc Natl Acad Sci USA* 96:10898–10901.
- Haskell CA, Cleary MD, and Chao IF (1999) Molecular understanding of fractalkine-mediated cell adhesion and signal transduction: Rapid flow arrest of CX3CR1-expressing cells is independent of G-protein activation. *J Biol Chem* 274:10053–10058.
- Hayashi S, Kurdoawa A, Miller EJ, Albright ME, Gorton BE and Cohen AB (1995) Synthetic bases and heptapeptides that inhibit IL-3 from binding to and activating human blood neutrophils. *J Immunol* 154:814–824.
- He J, Chen Y, Farzan M, Choe H, Ohagen A, Gartner S, Busciglio J, Yang X, Hoffmann W, Newman W, Mackay CR, Sodroski J and Gabuzda D (1997) CXCR3 and CXCR5 are co-receptors for HIV-1 infection of microglia. *Nature (Lond)* 385:645–649.
- Heath H, Qin S, Rao P, Wu L, LaRosa G, Kassam N, Ponath PD and Mackay CR (1997) Chemokine receptor function by human eosinophils: The importance of CXCR3 demonstrated using an antagonistic monoclonal antibody. *J Clin Invest* 99:178–184.
- Hebert CA, Vitangcol RV and Baker JB (1991) Scanning mutagenesis of interleukin-8 identifies a cluster of residues required for receptor binding. *J Biol Chem* 266:18989–18994.
- Hessom M, Bernier M, Ma, Hopken UE, Gerard NP and Dorf ME (1997) Alternative splicing of mouse fms/CXC chemokine receptor-4: stromal cell-derived factor-1alpha is a ligand for both CXC chemokine receptor-4 isoforms. *J Immunol* 158:3561–3564.
- Herbin G, Mahknecht U, Battilana F, Gregersen P, Pappas T, Butler J, O'Brien WA and Verdier E (1998) Apoptosis of CD8+ T cells is mediated by macrophages through interaction of HIV gp120 with chemokine receptor CXCR4. *Nature (Lond)* 395:189–194.
- Hesselgesse J, Chitnis CE, Miller LH, Yansura DG, Simmons LC, Fairbrother WJ, Kotts C, Wirth C, Gillet-Castro BL and Horak R (1993) A mutant of melanoma growth stimulating activity does not activate neutrophils but blocks erythrocyte invasion by malaria. *J Biol Chem* 268:11742–11745.
- Hesselgesse J, Halka-Miller M, DeVecchio V, Peiper SC, Hoxie J, Kolson DL, Taub D and Horak R (1997) CD4-independent association between HIV-1 gp120 and CXCR4: Functional chemokine receptors are expressed in human neurons. *Curr Biol* 7:112–121.
- Hesselgesse J, Ng HP, Liang M, Zheng W, May K, Bauman JG, Monahan S, Islam I, Wei GP, Ghannam A, Taub DD, Rosser M, Snider RM, Morrissey MM, Perez HD and Horak R (1998a) Identification and characterization of small molecule functional antagonists of the CCR1 chemokine receptor. *J Biol Chem* 273:15687–15692.
- Hesselgesse J, Taub D, Baakar P, Greenberg M, Hoxie J, Kolson DL and Horak R (1998b) Neuronal apoptosis induced by HIV-1 gp120 and the chemokine SDF-1alpha is mediated by the chemokine receptor CXCR4. *Curr Biol* 8:535–538.
- Heveker N, Montes M, Gernemroth L, Amara A, Trautman A, Alizon M and Schneider-Mergener J (1998) Dissociation of the signalling and antiviral properties of SDF-1-derived small peptides. *Curr Biol* 8:369–378.
- Holmes WE, Lee J, Kuang W, Rice GC and Wood WI (1991) Structure and functional expression of a human interleukin-8 receptor. *Science (Wash DC)* 253:1278–1280.
- Horney B, Wang W, Soto H, Buchanan M, Wiesenborn A, Catron D, Müller A, McClanahan T, O'Connor R, Ruzicki T, Lehmann P, Oltman E and Zlotnik A (2000) The orphan chemokine receptor GPR-2 (CCR10) binds the skin-associated chemokine CCL27 (CTACK/ALPILC). *J Immunol*, in press.
- Hoogwerf AJ, Black D, Proudfoot AEI, Wells TNC and Power CA (1996) Molecular cloning of murine CC CKR-4 and high affinity binding of chemokines to murine and human CC CKR-4. *Biochem Biophys Res Commun* 218:337–343.
- Horak R, Chitnis C, Darbonne W, Colby TJ, Rybicki A, Hadley TJ and Miller LH (1993) The erythrocyte chemokine receptor is a receptor for the malarial parasite *Plasmodium vivax*. *Science (Wash DC)* 261:1182–1184.
- Horak R (1994) The interleukin-8-receptor family: From chemokines to malaria. *Immunol Today* 15:169–174.
- Horak R, Hesselgesse J, Zhou Y, Fauds D, Halka-Miller M, Harvey S, Taub D, Samson M, Parmentier M, Rucker J, Doranz BJ and Doms RW (1998) The CC chemokine 1-309 inhibits CCR5-dependent infection by diverse HIV-1 strains. *J Biol Chem* 273:3986–3991.
- Horak R, Martin AW, Wang Z, Schweitzer L, Gerassimides A, Guo H, Lu Z, Hesselgesse J, Perez HD, Kim J, Parker J, Hadley TJ and Peiper SC (1997) Expression of chemokine receptors by subsets of neurons in the central nervous system. *J Immunol* 158:2828–2836.
- Howard OMZ, Openehm JG, Hollingshead MG, Covey JM, Bigelow J, McCormack JI, Buckheit RW, Clanton DJ, Turpin JA and Rice WG (1998) Inhibition of *in vitro* and *in vivo* HIV replication by a distancin analogue that interferes with chemokine receptor function: A candidate for chemotherapeutic and microbicidal application. *J Med Chem* 41:2184–2193.
- Huang Y, Paxton WA, Wolinsky SM, Neumann AJ, Zhang L, He T, Kang S, Ceradini D, Jin Z, Yazdanbakhsh K, Kunzmann K, Erickson D, Dragon E, Landau NR, Phair J, Ho DD and Koop RA (1996) The role of a mutant CXCR5 allele in HIV-1 transmission and disease progression. *Nat Med* 2:1240–1243.
- Huffnagle BG, McNeil LK, McDonald RA, Murphy JW, Towes BG, Maeda N and Kuziel WA (1999) Role of C-C chemokine receptor 5 in organ-specific and innate immunity to Cryptosporidium parvum. *J Immunol* 163:4642–4646.
- Humbles AE, Conroy DM, Marletta S, Raskin SM, Palfner RM, Proudfoot AEI, Wells TN, Li D, Jeffery RW, Ruzicki DA, Williams TJ and Jose PJ (1997) Kinetics of cotaxin generation and its relationship to eosinophil accumulation in allergic airway disease: Analysis in a guinea pig model *in vivo*. *J Exp Med* 186:601–612.
- Inai T, Bata M, Nishimura M, Kakizaki M, Takagi S and Yoshie O (1997a) The T cell-directed CC chemokine TARC is a highly specific biological ligand for CC chemokine receptor 4. *J Biol Chem* 272:15036–15042.
- Inai T, Chantry D, Raport JC, Ruzicki DA, Kunita M, Nishimura M, Godiska R, Yoshie O and Gray PW (1998) Macrophage-derived chemokine is a functional ligand for the CC chemokine receptor 4. *J Biol Chem* 273:1764–1768.
- Inai T, Hoshino K, Nishimura M, Kakizaki M, Bata M, Nishimura M, Kakizaki M, Takagi S, Nomiyama H, Schall TJ and Yoshie O (1997b) Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. *Cel* 91:521–530.
- Isewawa Y, Ping Z, Nakano K, Sugimoto N and Yamashiki K (1998) Human herpesvirus 6 open reading frame U12 encodes a functional beta-chemokine receptor. *J Virol* 72:6104–6112.
- Jen H, Cox MA, Kakizaki H, Zhang M, Byrnes H, Fine J, Lundell D, Chou CC, Narula SK and Zandovay PJ (1999) Cutting edge: species specificity of the CC chemokine receptor 4 (CXCR4) for stromal cell derived factor-1 (SDF-1). *J Immunol* 163:1269–1276.
- Johnston B, Burns AR, Suematsu M, Issekutz TB, Woodman RC and Kubers P (1999) Chronic inflammation upregulates chemokine receptors and induces neutrophil migration to monocyte chemoattractant protein-1. *J Cell Physiol* 108:1269–1276.
- Jones SA, Wolf M, Qin S, Mackay CR and Baggiolini M (1996) Different functions for the interleukin 8 receptors (IL-8R) of human neutrophil leukocytes: NADPH oxidase and phospholipase D are activated through IL-8R1 but not IL-8R2. *Proc Natl Acad Sci USA* 93:6682–6686.
- Jose PJ, Griffiths-Johnson DA, Collins PD, Walsh DT, Moschel R, Totty FN, Truong O, Hsuan JG and Williams TJ (1994) Eotaxin: A potent eosinophil chemoattractant cytokine defined in a guinea pig model of allergic airways inflammation. *J Exp Med* 179:851–856.
- Kaye D, Fagerström Wolf I, Ebersberger C, Kuehl WM and Lipp M (1993) The G protein-coupled receptor BLR1 is involved in murine B cell differentiation and is also expressed in neuronal tissues. *Eur J Immunol* 23:2532–2539.
- Kanbe K, Shimizu N, Suda Y, Takagishi K and Hoshino H (1999) A CXC chemokine receptor, CXCR4/BLR1, is a novel and specific receptor for human immunodeficiency virus type 2. *Virology* 265:264–273.
- Kellermann SA, Hudak S, Oldham ER, Liu YJ and McEvoy LM (1999) The CC chemokine receptor-7 (CCR7) binds CCR6 and macrophage inflammatory protein-3b but is not a chemokine receptor for *in vitro* and *in vivo* derived dendritic cells. *J Immunol* 162:3590–3594.
- Kelner GS, Kennedy J, Bacon BK, Kleyenstam S, Lagespades DA, Jenkins NA, Copeland NG, Bazan JF, Moore KW, Schall TJ and Zlotnik A (1994) Lymphotactin: A cytokine that represents a new class of chemokine. *Science (Wash DC)* 265:1935–1939.
- Kennedy KJ, Strieter RM, Kunkel SL, Lukacs NW and Karpus WJ (1998) Acute and relapsing experimental autoimmune encephalomyelitis are regulated by differential expression of the CC chemokines macrophage inflammatory protein-1alpha and monocyte chemoattractant protein-1. *J Neuroimmunol* 92:99–109.
- Kim H, Saito N, Inai T, Inai T, Takagi S, Sumi K, Nakajima S, Coughlin S, Tiffany HL, Murphy PM and Yoshie O (1996) Molecular cloning of human eotaxin, an eosinophil-selective CC chemokine, and identification of a specific eosinophil selectin receptor, CC chemokine receptor 3. *J Biol Chem* 271:725–730.
- Kim H, Saito N, Inai T, Takagi S, Sumi K, Nakajima S, Hirai K, Nomiyama H and Yoshie O (1999) Molecular cloning of a novel human CC chemokine (eotaxin-3) that is a functional ligand of CC chemokine receptor 3. *J Biol Chem* 274:27915–27920.
- Kitayama J, Mackay CR, Ponath PD and Springer TA (1998) The CC chemokine receptor CXCR3 participates in stimulation of eosinophil arrest on inflammatory endothelium in shear flow. *J Clin Invest* 101:2014–2017.
- Klodal TN, Rosenkilde MM and Schwartz TW (1998) Selective recognition of the membrane-bound CX3C chemokine, fractalkine, by the human cytomegalovirus-encoded broad spectrum receptor US28. *FEBS Lett* 441:209–214.
- Klodal TN, Rosenkilde MM, Coulin P, Simmons G, Johnsen AH, Alouani S, Power CA, Lütjeholm HK, Gerstoft J, Clapham PR, Clark-Lewis I, Wells TNC and Schwartz TW (1997) A broad spectrum chemokine antagonist: evidence for Kapozi's lymphoma-associated chemokine as a functional ligand for CXCR4. *Proc Natl Acad Sci USA* 94:2220–2225.
- Kowalska MA, Ratajczak J, Hoxie J, Brass LF, Gewirtz A, Poncz M and Ratajczak MZ (1999) Megakaryocyte precursors, megakaryocytes and platelets express the IL-8 receptor CXCR4 on their surface: Determination of response to stromal-derived factor-1 by megakaryocytes and platelets. *Br J Haematol* 104:220–225.
- Krutzik MD, Hromas R, Brown DR, Braxmeier UE and Fife KH (1997) Functional characterization of the C-C chemokine-like molecules encoded by *Moluccan contagionum* virus types 1 and 2. *Proc Natl Acad Sci USA* 94:9875–9880.
- Kuang Y, Wu Y, Jiang H and Wu D (1996) Selective G protein coupling by C-C chemokine receptors. *J Biol Chem* 271:15975–15978.
- Kurihara T and Bravo R (1996) Cloning and functional expression of mCCR2, a murine receptor for the C-C chemokines JE and FIC. *J Biol Chem* 271:11603–11607.
- Kurihara T, Warr G, Loy J and Bravo R (1997) Defects in macrophage recruitment and host defense in mice lacking the CXCR2 chemokine receptor. *J Exp Med* 186:1757–1762.
- Kuziel WA, Morgan SJ, Dawson TC, Griffin S, Smithies O, Ley K and Maeda N (1997) Severe reduction in leukocyte adhesion and monocyte extravasation in mice lacking the chemokine receptor 5. *Proc Natl Acad Sci USA* 94:12053–12058.
- LaCasse RA, Folis KE, Trahey M, Scarborough JD, Littman DR and Nunberg DH (1999) Fusion-competing vaccines: Broad neutralization of primary isolates of HIV. *Science (Wash DC)* 283:357–362.
- Laloni AS, Graham K, Moorman K, Rajarathnam K, Clark-Lewis I, Kelvin D and Mackay CR (1997) The partition of myxoma virus gamma interferon receptor homolog M-77 interacts with the heparin-binding domains of chemokines. *J Virol* 71:4356–4363.
- Laloni AS, Masters J, Zeng W, Barrett J, Pannu R, Everett H, Arendt CW and Mackay CR (1999) Usage of chemokine receptors by poxviruses. *Science (Wash DC)* 286:1968–1971.
- Laloni AS and McCadden G (1999) Evasion and exploitation of chemokines by viruses. *Cytokine Growth Factor Rev* 10:219–233.

- Lapham CK, Zaitseva MB, Lee S, Romanetsva T and Golding H (1999) Fusion of monocytes and macrophages with HIV-1 correlates with biochemical properties of CXCR4 and CCR5. *Nat Med* 5:633-638.
- Lee B, Doranz BJ, Rasmussen SA, Salas M, Frade JM, Martinez-A C, O'Brien SJ, Dean M, Collman RG and Doms RW (1998) Influence of the CCR2-V64I polymorphism on human immunodeficiency virus type 1 coreceptor activity and on chemokine receptor function of CCR2b, CCR3, CCR5, and CXCR4. *J Virol* 72:7450-7458.
- Lee J, Cicalano G, Camerato T, Toy K, Moore MW and Wood WI (1995) Chemokine binding and activities mediated by the mouse IL-8 receptor. *J Immunol* 155:2158-2164.
- Lee J, Horuk R, Rice GC, Bennett GL, Camerato T and Wood WI (1992) Characterization of two high affinity human interleukin-8 receptors. *J Biol Chem* 267:16255-16257.
- Legler DF, Loetscher M, Ross RS, Clark-Lewis I, Baggiolini M and Moser B (1998) B-cell-attracting chemokine 1, a human CXK chemokine expressed in lymphoid tissues, selectively attracts B lymphocytes via BLR/CXCR5. *J Exp Med* 187:655-660.
- Liao F, Lee HH and Farber JM (1997a) Cloning of STRL22, a new human gene encoding a G-protein-coupled receptor related to chemokine receptors and located on chromosome 6q27. *Genomics* 40:178-80.
- Liao F, Alderson K, Su J, Ulrich SJ, Kreider BL and Farber JM (1997b) STRL22 is a receptor for the CXK chemokine MIP-3alpha. *Biochem Biophys Res Commun* 236:212-217.
- Liao F, Alkhatib G, Peden KW, Sharma G, Berger EA and Farber JM (1997c) STRL33, a novel chemokine receptor-like protein, functions as a fusion cofactor for both macrophage-tropic and T cell line-tropic HIV-1. *J Exp Med* 185:2015-2023.
- Liao F, Rabin RL, Smith CS, Sharma G, Nutman TB and Farber JM (1999) CXK chemokine receptor 6 is expressed on diverse memory subsets of T cells and determines responsiveness to macrophage inflammatory protein 3 alpha. *J Immunol* 162:186-194.
- Libert F, Cochaux P, Beckman G, Samson M, Aksemsu N, Cao A, Czeizel A, Claustres M, de la Rua C, Ferrari M, Ferrec C, Glover G, Grinde B, Guaran S, Kucinasova V, Lavigne J, Merleir B, Ogar G, Peltonen I, Hostetler C, Schwartz M, Spitsyn V, Timar L, Beckman L, Vassart G and Parmentier M (1998) The delatase mutation conferring protection against HIV-1 in caucasian populations has a single and recent origin in northeastern Europe. *Hum Mol Genet* 7:399-406.
- Lindley LD, Westwick J and Kunkel SL (1993) Nomenclature announcement: The chemokines. *Immunol Today* 14:24.
- Ling K, Wang P, Zhao J, Wu YL, Cheng JJ, Xu GX, Hu W, Ma L and Pei G (1999) Five-transmembrane domains appear sufficient for a G-protein-coupled receptor: Functional five-transmembrane domain chemokine receptors. *Proc Natl Acad Sci USA* 96:7922-7927.
- Lira SA, Zaluska P, Heinrich JN, Fuentes ME, Carrasco D, Lewin AC, Barton DS, Durham S and Bravo R (1994) Expression of the chemokine N5/JKC in the thymus and epidemics of transgenic mice results in marked infiltration of a single class of inflammatory cells. *J Exp Med* 180:2039-2048.
- Liu R, Paxton WA, Choe S, Ceradini D, Martin SR, Horuk R, MacDonald ME, Stutman H, Koup RA and Landau NR (1996) Homologous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* 86:367-377.
- Loetscher P, Seitz M, Clark-Lewis I, Baggiolini M and Moser B (1994) Both interleukin-8 receptors independently mediate chemotaxis. Jurkat cells transfected with IL-8R1 or IL-8R2 migrate in response to IL-8, GRO alpha and NAP-2. *FEBS Lett* 341:187-192.
- Loetscher M, Amara A, Oberlin E, Brass N, Legler D, Loetscher P, D'Apuoso M, Meese R, Rousset D, Virelizier JH, Baggiolini M, Arenzana-Seisdedos F and Moser B (1997) TYMSRTR, a putative chemokine receptor selectively expressed in activated T cells, exhibits HIV-1 coreceptor function. *Curr Biol* 7:652-660.
- Loetscher M, Gerber B, Loetscher P, Jones SA, Piali L, Clark-Lewis I, Baggiolini M and Moser B (1996) Chemokine receptor specific for IP10 and Mig: Structure, function, and expression in T lymphocytes. *J Exp Med* 184:963-970.
- Loetscher M, Loetscher P, Brass N, Meese E and Moser B (1998a) Lymphocyte-specific chemokine receptor CXCR3: Regulation, chemokine binding and gene localization. *Eur J Immunol* 28:3696-3705.
- Loetscher P, Gong JH, Dewald B, Baggiolini M and Clark-Lewis I (1998b) N-terminal peptides of stromal cell derived factor-1 with CXK chemokine receptor 4 agonist and antagonist activities. *J Biol Chem* 273:22279-22283.
- Loetscher P, Seitz M, Baggiolini M and Moser B (1996) Interleukin-2 regulates CXK chemokine receptor expression and chemotactic responsiveness in T lymphocytes. *J Exp Med* 184:569-577.
- Loetscher P, Uguccioni M, Bordoli L, Baggiolini M, Moser B, Chizzolini and Dayer JM (1998c) CCR5 is characteristic of Th1 lymphocytes. *Nature (Lond)* 391:344-345.
- Lomize AL, Pogorelec ID and Mosberg HL (1999) Structural organization of G-protein-coupled receptors. *J Computat Mol Biol* 12:325-353.
- Ludwig A, Petersen F, Zahn S, Gotze O, Schroder JM, Flad HD and Brandt E (1997) The CXK chemokine neutrophil-activating peptide-2 induces two distinct options of neutrophil chemotaxis by differential interaction with interleukin-5 receptors CXCR-1 and CXCR-2. *Blood* 89:4588-4597.
- Luo H, Chaudhuri A, Johnson KR, Neote K, Zhelezna Y, He Y and Poppo AO (1997) Cloning, characterization, and mapping of a murine promiscuous chemokine receptor gene: Homolog of the human Duffy gene. *Genome Res* 7:932-941.
- Luster AD (1998) Chemokines: Chemokinetic cytokines that mediate inflammation. *N Engl J Med* 338:458-463.
- Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T, Bronson RT and Springer TA (1998) Impaired B-lymphopoiesis, myelopoiesis, and derelict cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci USA* 95:9448-9453.
- Mallatou G, Soo KS, Schall TJ, Pisarska M and Anstee DJ (1995) Mutations in the erythrocyte chemokine receptor (Duffy) gene: The molecular basis of the Fya/Fyb antigens and identification of a deletion in the Duffy gene of an apparently healthy individual with the Fy(a-b-) phenotype. *Br J Haematol* 90:823-829.
- Marchese A, Decheray JM, Nguyen T, Heng HX, Salas M, Cheng R, Murphy PM, Tsou LC, Shi X, Greer P and O'Dowd B (1995) Cloning of human genes encoding novel G-protein-coupled receptors. *Genomics* 23:609-618.
- Marshall E, Howell AH, Powles R, Hunter MG, Edwards M, Wood LM, Czaplewski L, Puttick R, Warrington S, Boyer M, Tusta N, Dexter TM, Lord BI and Millar A (1998) Clinical effects of human macrophage inflammatory protein-1 alpha MIP-1 alpha (LD78) administration to human: A phase I study in cancer patients and normal healthy volunteers with the genetically engineered variant, BB-10010. *Eur J Cancer* 34:1023-1029.
- Martin MP, Dean M, Smith MW, Winkler C, Gernard R, Michael NL, Lee B, Doms RW, Margalit J, Bauminger S, Goed JJ, O'Brien TR, Hargitator MW, Vlahov D, O'Brien SJ and Carrington M (1998) Genetic acceleration of AIDS progression by a promoter variant of CCR5. *Science* 282:1907-1911.
- Massung RF, Jayarama V and Meyer RW (1993) DNA sequence analysis of conserved and unique regions of swinepox virus: Identification of protein-coding sequences supporting phenotypic observations including a novel G-protein-coupled receptor homologue. *Virology* 197:511-528.
- Matsumura K, Larsen CG, Dubois GO and Oppenheim JJ (1989) Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line. *J Exp Med* 169:1455-1490.
- Mathew AN, Lind DS, Zimmermann N, Saraf MN, Luster AD, Pearlman E, Wert SE and Rothberg ME (1998) Eotaxin is required for the baseline level of tissue eosinophilia. *Proc Natl Acad Sci USA* 95:6273-6278.
- McMordet HD, Zimmermann PA, Guignard F, Kleberberg CA, Leliman SF and Murphy PM (1998) CCR5 promoter polymorphism and HIV-1 disease progression. *Lancet* 352:896-870.
- Mellado M, Rodriguez-Frade JM, Vila-Coro AJ, de Ana AM and Martinez-A C (1999) Chemokine control of HIV-1 infection. *Nature (Lond)* 400:723-724.
- Meucci O, Fatate A, Simon AN, Bushnell TJ, Gray RW and Miller RJ (1998) Chemokines regulate hippocampal neuronal signaling and gp120 neurotoxicity. *Proc Natl Acad Sci USA* 95:14500-14505.
- Meyer A, Coyle AJ, Proudfoot AE, Wells TN and Power CA (1996) Cloning and characterization of a novel murine macrophage inflammatory protein-1 alpha homologue. *Genes Dev* 10:2126-2131.
- Michael NL, Nelson JA, Kewal Ramani VN, Chang G, O'Brien SJ, Mascola JR, Volsky B, Louder M, White OC 2nd, Littman DR, Swanstrom R and O'Brien TR (1996) Exclusive and persistent use of the entry coreceptor CXCR4 by human immunodeficiency virus type 1 from a subject homozygous for CCR5 delta32. *J Virol* 71:4445-4451.
- Miller LH, Mason SJ, Clyde DF and McGinniss MH (1976) The resistance factor to *Plasmodium vivax* in blacks: The Duffy-blood-group genotype, Fy(a-b-). *J Med Res* 155:455-462.
- Miller LH, Mason SJ, Dorosh J, McGinniss MH and Rothman RI (1975) Erythrocyte receptor for *Plasmodium knowlesi* malaria: Duffy blood group determinants. *Science (Wash DC)* 189:561-563.
- Modi VS and Yoshimura T (1999) Isolation of novel GRO genes and a phylogenetic analysis of the CXK chemokine subfamily. *Mol Biol Cell* 10:189-193.
- Montecarlo FS and Charo IF (1995) The amino-terminal extracellular domain of the MCP-1 receptor, but not the RANTES/MIP-1alpha receptor, confers chemokine selectivity: Evidence for a two-step mechanism for MCP-1 receptor activation. *J Biol Chem* 271:19984-19992.
- Moore MW, Cicalano G and Wood WI (1995) Technical comment. *Science (Wash DC)* 269:1591.
- Morales J, Horney B, Vicari AP, Hudak S, Oldham E, Hadrack J, Orozco R, Copeland NG, Jenkins NA, McEvoy LM and Zlotnik A (1999) CTACK, a skin-associated chemokine that preferentially attracts skin-homing memory T cells. *Proc Natl Acad Sci USA* 96:14470-14475.
- Morohashi H, Miyawaki T, Nomura H, Kuno K, Murakami S, Matsushima K and Mukaida N (1995) Expression of both types of human interleukin-8 receptors on mature neutrophils, monocytes, and natural killer cells. *J Leukoc Biol* 57:189-187.
- Moser B, Schumacher C, von Tiechauer V, Clark-Lewis I and Baggiolini M (1991) Neutrophil-activating peptide 2 and granuloma-oncogene growth-stimulatory activity interact with neutrophil-activating peptide/interleukin 8 receptors on human neutrophils. *J Biol Chem* 266:10660-10671.
- Murakami T, Nakajima T, Koyanagi Y, Tachibana K, Fujii N, Tamamura H, Yoshida A, Waki M, Matsumoto A, Yoshida C, Kishimoto T, Yamamoto N and Nagasawa T (1997) A small molecule CXCR4 inhibitor that blocks T cell line-tropic HIV-1 infection. *J Exp Med* 186:1389-1393.
- Murphy PM (1993) Molecular mimicry and the generation of host defense protein diversity. *Cell* 72:817-826.
- Murphy PM (1994) The molecular biology of leukocyte chemoattractant receptors. *Annu Rev Immunol* 12:593-633.
- Murphy PM (1996) Chemokine receptors: Cloning strategies. *METHODS: A Companion to Methods in Enzymology* 10:104-115.
- Murphy PM and Littman HL (1991) Cloning of complementary DNA encoding a functional interleukin-8 receptor. *Science* 253:1280-1283.
- Myers SJ, Wong LM and Charo IF (1995) Signal transduction and ligand specificity of the human monocyte chemoattractant protein-1 receptor in transfected embryonic kidney cells. *J Biol Chem* 270:5766-5774.
- Nagasawa T, Hirota S, Tachibana K, Takikawa N, Nishikawa S, Kitamura Y, Yoshida N, Kikutani H and Kishimoto T (1996) Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXK chemokine PBSF/SDF-1. *Nature (Lond)* 382:635-638.
- Najakshani AM, Mochizuki LV, Alayash VI and Taranin AV (1999) Identification of an IL-8 homologue in lamprey (*Lampetra fluviatilis*): Early evolutionary divergence of chemokines. *Eur J Immunol* 29:375-382.
- Napolitano M, Zingoni A, Bernardini G, Spinetti G, Nista A, Storlazzi CT, Rocchi M

- and Santoni A (1996) Molecular cloning of TER1, a chemokine receptor-like gene expressed by lymphoid tissues. *J Immunol* 157:3729–3763.
- Nardelli B, Tiffany HL, Bong DW, Yourey PA, Morahan DR, Li Y, Murphy PM and Alderson RF (1999) Characterization of the signal transduction pathway activated in human monocytes and dendritic cells by MIP1- α , a specific ligand for CC chemokine receptor 1. *J Biol Chem* 274:4354–444.
- Neote K, DiGregorio D, Mak JY, Honk R and Schall TJ (1993) Molecular cloning, functional expression, and signaling characteristics of a C-C chemokine receptor. *Cell* 72:415–428.
- Neote K, Mak JY, Kolakowski LF Jr and Schall TJ (1994) Functional and biochemical analysis of the cloned Duffy antigen: Identity with the red blood cell chemokine receptor. *Blood* 84:44–52.
- Ng HF, May K, Bauman JG, Ghannam A, Isen I, Liang M, Horuk R, Hesselgesell J, Snider RM, Perez HD and Morrissey JM (1999) Discovery of novel non-peptide CCR1 receptor antagonists. *J Med Chem* 42:4680–4694.
- Nibbs RJ, Salcedo TW, Campbell JD, Yao XT, Li Y, Nardelli B, Olsen HS, Morris TS, Proudfoot AE, Patel PV and Graham GJ (2000) C-C chemokine receptor 3 antagonism by the β -chemokine macrophage inflammatory protein 4, a property strongly enhanced by an amino-terminal alanine-methionine swap. *J Immunol* 164:1488–1497.
- Nibbs RJ, Wylie SM, Yang J, Landau NR and Graham GJ (1997b) Cloning and characterization of a novel promiscuous human beta-chemokine receptor D6. *J Biol Chem* 272:32078–32083.
- Nibbs RJ, Wylie SM, Pragnell IB and Graham GJ (1997a) Cloning and characterization of a novel murine beta chemokine receptor, D6: Comparison to three other related macrophage inflammatory protein-1alpha receptors. *CCR1*, CCR3, and CCR5. *J Biol Chem* 272:14858–14864.
- Nibbs RJ, Yang J, Landau NR, Mao JH and Graham GJ (1999) LD78beta, a non-allelic variant of human MIP-1alpha (LD78alpha), has enhanced receptor interactions and potent HIV suppressive activity. *J Biol Chem* 274:17478–17483.
- Nicholas J (1996) Determination and analysis of the complete nucleotide sequence of human herpesvirus 7. *J Virol* 70:5975–5989.
- Nilsen G, Mikovits JA, Metcalfe DD and Taub DD (1999) Mast cell migratory response to interleukin-8 is mediated through interaction with chemokine receptor CXCR2/interleukin-8RB. *Blood* 93:2791–2797.
- Oberlin E, Amara A, Bachelier F, Besia C, Virelizier JL, Arenzana-Seisdedos F, Schwartz O, Huet JM, Clark-Lewis I, Legler DF, Loetscher M, Baggiolini M and Moser B (1996) The CXCR chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-adapted HIV-1. *Nature (Lond)* 382:833–835.
- Oehminger B, Tassera L, Biffare D, Rhs S and Dahinden CA (1999) Regulation of cytokine expression and chemokine formation in human basophils by growth factors, chemokines and chemotactic agonists. *Eur J Immunol* 29:11–22.
- Ochi H, Hirani WM, Yuan Q, Friend DS, Austen KF and Boyce JA (1999) T helper cell type 2 cytokine-mediated chemotactic responses and CCR3 expression during differentiation of human mast cells in vitro. *J Exp Med* 190:267–290.
- Olsen WC, Rabut CR, Nagashima KA, Tran DN, Almeida DJ, Mosier SP, Segal JP, Thompson DA, Kajumo F, Guo Y, Moore JP, Madden JP and Dragic T (1999) Differential inhibition of human immunodeficiency virus type 1 fusion, gp120 binding, and CC-chemokine activity by monoclonal antibodies to CCR5. *J Virol* 73:1145–1155.
- Oppenheim JJ, Zachariae CO, Mukaida N and Matsushima K (1991) Properties of the novel proinflammatory supergene "interleukin" cytokine family. *Annu Rev Immunol* 9:617–648.
- Oppenheim JJ, Howard OMT and Gotsel E (2000) *The Role of Chemotactic Factors and Neuropeptides in Host Defense*. Online Cytokine Textbook. Academic Press, San Diego.
- Oravecz T, Pall M, Rodriguez G, Correll MD, Ditto M, Nguyen NY, Boykins R, Unsworth E and Norcross MA (1997) Regulation of the receptor specificity and function of the chemokine RANTES (regulated on activation, normal T cell expressed and secreted) by dipeptidyl peptidase IV (CD26)-mediated cleavage. *J Exp Med* 186:1865–1872.
- Owman C, Garzino-Demo A, Cocchi F, Popovic M, Sabharwal A and Gallo RC (1998) The leukotriene B₄ receptor functions as a novel type of chemotactic mediating entry of primary HIV-1 isolates into CD4-positive cells. *Proc Natl Acad Sci USA* 95:9530–9534.
- Pal R, Garzino-Demo A, Markham PD, Burns J, Brown M, Gallo RC and DeVico AL (1997) Inhibition of HIV-1 infection by the beta-chemokine MDC. *Science (Wash DC)* 278:695–698.
- Palframan RT, Collins PD, Williams TJ and Rankin SM (1998) Estrogen induces a rapid release of eosinophils and their progenitors from the bone marrow. *Blood* 91:2240–2248.
- Pan L, Dussault B, Woolf E, Alperin G, Culpepper J, Gutierrez-Ramos JC, Gearing D, Lian Y, Lloyd C, Zhou H, Dolich S, Deeds J, Gonzalo JA, Vith J, Gosselin M and Ma J (1997) Neutrophils, a subset of chemokine-anchored chemokine upregulated in brain inflammation. *Nature (Lond)* 387:611–617 [erratum published in *Nature (Lond)* 1997:388:100].
- Pease JE and Murphy PM (1998) Microbial corruption of the chemokine system: an expanding paradigm. *Semin Immunol* 10:169–178.
- Pease JE, Wang J, Posaht PD and Murphy PM (1998) The N-terminal extracellular segments of the chemokine receptors CCR1 and CCR3 are determinants for MIP-1alpha and chemokine binding, respectively, but a second domain is essential for efficient receptor activation. *J Biol Chem* 273:19772–19776.
- Pfeiffer SC, Wang ZX and Neufeldt A (1999) The Duffy negative phenotype for chemokines (DARC) is expressed in endothelial cells of Duffy negative individuals who lack the erythrocyte receptor. *J Exp Med* 191:1311–1317.
- Poled A, Petit I, Kollet O, Magid M, Ponomarev T, Byk T, Nagler A, Ben-Hur H, Many A, Shultz L, Lidor O, Alon R, Zipori D and Lapidot T (1999) Dependence of human stem cell engraftment and repopulation of NODSCID mice on CXCR4. *Science (Wash DC)* 283:845–848.
- Penfold MET, Dairaghi DJ, Duke GM, Saeedur N, Mocarski ES, Kemble GW and Schall TJ (1999) Cytomegalovirus encodes a potent alpha chemokine. *Proc Natl Acad Sci USA* 96:9339–9344.
- Petering H, Gotoe O, Kimmig D, Smolarski R, Kapp A and Eisner J (1999) The chemokine receptor CXCR1: Functional analysis and expression of CXCR1 and CXCR2 on human eosinophils. *Blood* 83:684–702.
- Pleskoff O, Treboute C and Alizon M (1998) The cytomegalovirus-encoded chemokine receptor US28 can enhance cell-cell fusion mediated by different viral proteins. *J Virol* 72:6389–6394.
- Pleskoff O, Treboute C, Brelet A, Heveker N, Seman M and Alizon M (1997) Identification of a chemokine receptor encoded by human cytomegalovirus as a cofactor for HIV-1 entry. *Science (Wash DC)* 276:1874–1878.
- Ponath PD, Qin S, Ringler I, Clark-Lewis I, Wang J, Kassam N, Smith H, Jia G-Q, Heuser M, Li Y, Gutierrez-Ramos JC and Mackay CR (1996) Cloning of the human eosinophil chemoattractant, eotaxin: Expression, receptor binding and functional properties provide a mechanism for the selective recruitment of eosinophils. *J Clin Invest* 97:604–612.
- Post TW, Bronic CR, Rothenberg ME, Luster AD, Gerard NP and Gerard C (1995) Molecular characterization of two murine eosinophil β chemokine receptors. *J Immunol* 155:5299–5305.
- Power CA, Clementson JM, Clementson KJ and Wells TN (1995a) Chemokine and chemokine receptor mRNA expression in human platelets. *Cytokine* 7:479–482.
- Power CA, Church DJ, Meyer A, Alouani S, Proudfoot AE, Clark-Lewis I, Sotgiu S, Mantovani A and Wells TN (1997) Cloning and characterization of a specific receptor for the novel CC chemokine MIP-1alpha from lung dendritic cells. *J Exp Med* 186:825–835.
- Power CA, Meyer A, Nemeth K, Bacon BK, Hoogwerf AJ, Proudfoot AE and Wells TN (1995b) Molecular cloning and functional expression of a novel CC chemokine receptor cDNA from a human basophilic cell line. *J Biol Chem* 270:19495–19500.
- Premack AB and Schall TJ (1996) Chemokine receptors: Gateways to inflammation and infection. *Nat Med* 2:1174–1178.
- Proudfoot AE, Power CA, Hoogwerf AJ, Montejano MO, Borlat F, Offord RE and Wells TN (1996) Extension of recombinant human RANTES by the retention of the initiating methionine produces a potent antagonist. *J Biol Chem* 271:2599–2603.
- Qin S, Rottman JB, Myers P, Kassam N, Weinblatt M, Loetscher M, Koch AE, Moser B and Mackay CR (1998) The chemokine receptors CXCR3 and CCR5 mark subsets of T cells associated with certain inflammatory reactions. *J Clin Invest* 101:746–754.
- Rabin RL, Park MK, Liao P, Swafford R, Stephany D and Farber JM (1999) Chemokine receptor responses on T cells are achieved through regulation of both receptor expression and signaling. *J Immunol* 162:3840–3850.
- Rauschanskas CM (1998) Chemokine involvement in MS tissue: Adhesion molecules and chemokines. *J Neuroimmunol* 89:57–68.
- Raport CJ, Gossling J, Schwelckart VL, Gray PW and Charo IF (1996) Molecular cloning and functional characterization of a novel human CC chemokine receptor (CCR5) for RANTES, MIP-1 α , and MIP-1 β . *J Biol Chem* 271:17161–17166.
- Reich Z, Gearing D, Roring L, Gossling J, Gearing A, Charo IF and Broxmeyer HE (1999) Enhanced myeloid progenitor cell cycling and apoptosis in mice lacking the chemokine receptor, CCR2. *Blood* 93:1524–1533.
- Rizazadeh CD, Wyatt R, Hernandez-Ramos N, Sun Y, Kwong PD, Hendrickson WA and Wells TN (1998) HIV-1: A conserved HIV-1 glycoprotein structure involved in chemokine receptor binding. *Science (Wash DC)* 280:1949–1953.
- Rodriguez-Frade JM, Vila-Coro AJ, de Ana AM, Albar JP, Martinez-A C and Mellado M (1999) The chemokine monocyte chemoattractant protein-1 induces functional responses through dimerization of its receptor CCR2. *Proc Natl Acad Sci USA* 96:3628–3633.
- Roos RS, Loetscher M, Legler DF, Clark-Lewis I, Baggiolini M and Moser B (1997) Identification of CCR3, the receptor for the human CC chemokine I-309. *J Biol Chem* 272:17251–17254.
- Rosenkilde MM, Kiedal TN, Branner-Osburne H and Schwartz TW (1990) Agonists and inverse agonists for the herpesvirus 8-encoded constitutively active seven-transmembrane oncogene product, ORF74. *J Biol Chem* 274:9586–9591.
- Rothenberg ME, MacLean JA, Pearlman E, Luster AD and Leder P (1997) Targeted disruption of the chemokine receptor partially reduces antigen-induced tissue eosinophilia. *J Exp Med* 185:785–790.
- Rubbert A, Combadiere C, Ostrowski M, Arthos J, Dwyer M, Machado E, Cohn MA, Hoxie JA, Murphy PM, Faud AS and Weissman D (1998) Dendritic cells express multiple chemokine receptors used as coreceptors for HIV entry. *J Immunol* 160:3928–3941.
- Rueda J, Gossling AL, Sharon M, Samson M, Lee B, Person JF, Yi Y, Margulies B, Collman RG, Doranz BJ, Parmentier M and Doms RW (1997) Utilization of chemokine receptors, orphan receptors, and herpesvirus-encoded receptors by diverse human and simian immunodeficiency viruses. *J Virol* 71:8999–9007.
- Rutter MW, Clark A, Artias J, Kinnaird A, Gattmair AT, Adelsberger J, Holmes KL, Cohen CB and Ransley MJ (1998) Peripheral blood-derived CD34⁺ progenitor cells: CXCR chemokine receptor 4 and CC chemokine receptor 5 expression and infection by HIV. *J Immunol* 161:4169–4176.
- Sabroe I, Conroy DM, Gerard NP, Li Y, Collins PD, Post TW, Jose PJ, Williams TJ, Giers GJ and Posaht PD (1998) Cloning and characterization of the guinea pig eosinophil eotaxin receptor, C-C chemokine receptor 3: Blockade using a monoclonal antibody in vivo. *J Immunol* 161:6139–6147.
- Saeedur N, Lin YC, Dairaghi DJ, Schall TJ and Mocarski ES (1999) Cytomegalovirus-induced beta chemokine promotes monocyte-associated viremia in the host. *Am J Pathol* 156:1081–1086.
- Saeki H, Moore AM, Brown MJ and Hwang ST (1999) Secondary lymphoid-tissue chemokine (SLC) and CC chemokine receptor 7 (CCR7) participate in the emigration pathway of mature dendritic cells from the skin to regional lymph nodes. *J Immunol* 162:2472–2475.
- Saleh M, Wasserman K, Young HA, Grimm MC, Howard OM, Anver MR, Kleinman HK, Murphy JD and Oppenheim JJ (1999) Vascular endothelial growth factor and basic fibroblast growth factor induce expression of CXCR4 on human endothelial

- cells: In vivo neovascularization induced by stromal-derived factor-1alpha. *Am J Pathol* 154:1125–1135.
- Sallusto F, Palermo B, Hoy A and Lanzavecchia A (1999a) The role of chemokine receptors in directing traffic of naive, type 1 and type 2 T cells. *Curr Top Microbiol Immunol* 246:123–129.
- Sallusto F, Kremmer E, Palermo B, Hoy A, Ponzetti P, Qin S, Forster R, Lipp M and Lanzavecchia A (1999b) Switch in chemokine receptor expression upon TCR stimulation reveals novel homing potential for recently activated T cells. *Eur J Immunol* 29:2037–2045.
- Sallusto F, Lenig D, Mackay CR and Lanzavecchia A (1998) Flexible programs of chemokine receptor expression on human polarized T helper 1 and 2 lymphocytes. *J Exp Med* 187:875–883.
- Sallusto F, Mackay CR and Lanzavecchia A (1997) Selective expression of the eotaxin receptor CCR3 by human T helper 2 cells. *Science (Washington DC)* 277:2005–2007.
- Sallusto F, Lenig D, Forster R, Lipp M and Lanzavecchia A (1999) Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature (London)* 401:708–712.
- Samson M, Edinger AL, Stordeur P, Rucker J, Verhaselt V, Sharroun M, Govaerts C, Molleuex C, Vassart G, Doms RW and Parmentier M (1996) ChemR23, a putative chemotactant receptor, is expressed in monocyte-derived dendritic cells and macrophages and is a coreceptor for HIV and some primary HIV-1 strains. *Eur J Immunol* 26:1689–1700.
- Samson M, Labbe O, Molleuex C, Vassart G and Parmentier M (1996a) Molecular cloning and functional expression of a new human CC-chemokine receptor gene. *Biochemistry* 35:6567–6573.
- Samson M, Libert F, Doranz BJ, Rucker J, Lisnard C, Farber CM, Saragosti S, Lapomerolles C, Cogniaux J, Forcellec C, Muyldermans G, Verhofstede C, Burton G, Georges M, Imal T, Rana S, Yi Y, Smyth RJ, Collman RG, Doms RW, Vassart G and Parmentier M (1996b) Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. *Nature (London)* 382:722–725.
- Samson M, Soularue P, Vassart G and Parmentier M (1996c) The gene encoding the human CC-chemokine receptors CC-CKR1 to CC-CKR5 (CMK1R1 to CMK5R5) are clustered in the p21-22/q24 region of chromosome 3. *Genomics* 30:522–526.
- Sarraf MN, Garcia-Zepeda EA, MacLean JA, Charo IF and Luster AD (1997) Murine monocyte chemotactic protein (MCP)-5: A novel CC chemokine that is a structural and functional homologue of human MCP-1. *J Exp Med* 185:899–1009.
- Schall TJ (1991) Biology of the RANTES/SIS cytokine family. *Cytokine* 3:165–183.
- Schall TJ, Bacon K, Toy KD and Goeddel DV (1990) Selective attraction of monocytes and T lymphocytes of the memory phenotype by cytokine RANTES. *Nature (London)* 347:669–671.
- Schols D, Struyf S, Van Damme J, Este JA, Henson G and De Clercq E (1997) Inhibition of HIV-1 HIV strains by selective antagonization of the chemokine receptor CXCR4. *J Exp Med* 186:1353–1358.
- Schwarz MK and Wells TM (1999) Interfering with chemokine networks: The hope for new therapeutics. *Curr Opin Chem Biol* 3:417–418.
- Schweickart VL, Epp A, Raport C and Gray PW (1994b) CCRL1 is a functional receptor for the MCP family of chemokines. *J Biol Chem*, in press.
- Schweickart VL, Raport CJ, Godlaski R, Byers MG, Eddy RL Jr, Shows TB and Gray PW (1994) Cloning of human monocyte EB1, a lymphoid-specific G-protein-coupled receptor encoded on human chromosome 17q12–q21.6. *Genomics* 23:843–850.
- Sekido N, Mukaida N, Harada A, Nakanishi I, Watanabe Y and Matsushima K (1993) Prevention of lung reperfusion injury in rabbits by a monoclonal antibody against interleukin-8. *Nature (London)* 365:654–657.
- Shimizu N, Soda Y, Kanbe K, Liu HY, Mukai R, Kitamura T and Hoshino H (2000) A putative G protein-coupled receptor, RD1C1, is a novel coreceptor for human and simian immunodeficiency viruses. *J Virol* 74:619–626.
- Shinkai A, Yoshida H, Koike M, Shoji E, Nakagawa S, Saito A, Takeda T, Imabeppu S, Kato Y, Hanae N, Aizawa H, Kuga T and Nishi T (1999) A novel human CC chemokine, eotaxin-3, which is expressed in IL-6-stimulated vascular endothelial cells, exhibits potent activity toward eosinophils. *J Immunol* 163:1602–1610.
- Shizura M, Nakano T, Inazawa J, Tashiro K, Tada H, Shinohara T and Honjo T (1996) Structure and chromosomal localization of the human stromal cell-derived factor 1 (SDF1) gene. *Genomics* 30:495–500.
- Simmons G, Clapham PR, Picard L, Offord RE, Rosenkilde MN, Schwartz TW, Buser R, Wells TNC and Proudfoot AE (1997) Potent inhibition of HIV-1 infectivity in macrophages and lymphocytes by a novel CCR5 antagonist. *Science (Washington DC)* 276:276–279.
- Smith CA, Smith TD and Goodwin RG (1997) Tlx porovirus genome encodes a secreted, soluble protein that preferentially inhibits type B chemokine activity yet lacks selectivity to known chemokine receptors. *Virology* 236:316–322.
- Smith MW, Dean M, Carrington M, Winkler C, Hootsley GA, Lomb DA, Goeken J, O'Brien TK, Jacobs-Lindner S, Buchbinder S, Vittinghoff E, Vlahov D, Hoek K, Hargarten MW, Hemophilia Growth and Development Study (HGDS), Multicenter AIDS Cohort Study (MACS), Multicenter Hemophilia Cohort Study (MHCS), San Francisco City Cohort (SFCC), ALIVE Study and O'Brien SJ (1997b) Contrasting genetic influences of CCR5 and CCR2 variants on HIV-1 infection and disease progression. *Science (Washington DC)* 277:859–864.
- Soto H, Wang W, Strieter RM, Copeland NG, Gilbert DM, Jenkins NA, Hedrick J and Zlotnik A (1998) The CC chemokine 6CKine binds the CX3C chemokine receptor CXCR3. *Proc Natl Acad Sci USA* 95:8250–8251.
- Sozzani S, Allavena P, D'Amico G, Vercelli A, Bianchi G, Katsura M, Imai T, Yoshie O, Bonacchi R and Mantovani A (1998a) Differential regulation of chemokine receptors during cellular maturation: A model for their trafficking properties. *J Immunol* 161:1083–1096.
- Sozzani S, Allavena P, Vercelli A and Mantovani A (1999) The role of chemokines in the regulation of dendritic cell trafficking. *J Leukoc Biol* 66:1–9.
- Sozzani S, Linali W, Bianchi G, Allavena P, Wells TM, Napolitano M, Bernardini G, Vercelli A, D'Amico G, Mazzeo D, Sinigaglia F, Santoni A, Maggi E, Romagnani G and Mantovani A (1998b) The viral chemokine chemokine inflammatory protein-II is a selective Th2 chemotactant. *Blood* 92:4038–4039.
- Sozzani S, Linali W, Bonacchi A, Polastri A, Nicosi L, Ziosi D, Piemonti L, D'Amico G, Power CA, Wells TM, Gobbi M, Allavena P and Mantovani A (1997) Receptor expression and responsiveness of human dendritic cells to a defined set of CC and CX3C chemokines. *J Immunol* 159:1993–2000.
- Springer TA (1994) Traffic signals for lymphocyte recirculation and leukocyte emigration: The multistep paradigm. *Cell* 76:301–314.
- Strebholz DN, Soderberg-Nauler C, Vieira J, Smith P, Wakabayashi E, Ruchti F, Mattsson K, Altschuler Y and Nelson JA (1999) The human cytomegalovirus chemokine receptor US28 modulates vascular smooth muscle cell migration. *Cell* 99:511–521.
- Struyf S, De Meester I, Schepers S, Lannetys JP, Menten P, Wang JM, Proost P and Van Damme J (1998) Natural truncation of RANTES abolishes signaling through the CC chemokine receptors CCR1 and CCR3, impairs its chemotactic potency and generates a chemokine inhibitor. *Eur J Immunol* 28:1262–1271.
- Su S, Mukaida N, Wang J, Zhang Y, Takami A, Nakano T and Matsushima K (1997) Inhibition of immature erythroid progenitor cell proliferation by macrophage inflammatory protein-1alpha by interacting mainly with a C-C chemokine receptor, CCR1. *Blood* 90:605–611.
- Su SB, Mukaida N, Wang J, Nomura H and Matsushima K (1996) Preparation of specific polyclonal antibodies to a C-C chemokine receptor, CCR1, and determination of CCR1 expression on various types of leukocytes. *J Leukoc Biol* 60:658–666.
- Suzuki G, Sawa H, Kobayashi Y, Nakata Y, Nakagawa K, Uzawa A, Sakiyama H, Kakimura S, Iwabuchi K and Nagashima K (1999) Pertussis toxin-sensitive signal transduction in the regulation of thymocyte survival at the corticosterone junction in the thymus. *J Immunol* 162:5981–5985.
- Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, Kitamura Y, Matsushima K, Yoshida N, Nishikawa S, Kishimoto T and Nagasawa T (1998) The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature (London)* 393:591–594.
- Tamamura H, Yu Y, Hattori T, Zhang X, Arakaki R, Kanbara K, Omagari A, Otsuka A, Iwuka T, Xiamamoto N, Nakashima H and Fujii N (1998) A low-molecular-weight inhibitor against the chemokine receptor CXCR4: A strong anti-HIV peptide. *T160. Biochem Biophys Res Commun* 253:787–793.
- Tanaka Y, Adams DH and Shaw S (1993) Proteoglycans on endothelial cells present adhesion-inducing cytokines to leukocytes. *Immunol Today* 14:111–115.
- Tang HL and Cyster JG (1999) Chemokine up-regulation and activated T cell attraction by myeloid dendritic cells. *Science (Washington DC)* 284:819–822.
- Tang T, Owen JD, Du J, Walker CL and Richmond A (1998) Molecular cloning and characterization of a mouse gene with homology to the Duffy-antigen receptor for chemokines. *DNA Seq* 9:120–133.
- Tashiro K, Nakamura T and Honjo T (1999) The signal sequence trap method. *Mol Cell* 13:473–485.
- Tashiro K, Tada H, Heliker R, Shirouzu M, Nakano T and Honjo T (1993) Signal sequence trap: A cloning strategy for secreted proteins and type I membrane proteins. *Science (Washington DC)* 261:600–603.
- Taylor TH and Hogenkamp C (1993) The DNA sequence of the DNA sequence of equine herpesvirus 2. *J Mol Biol* 234:520–529.
- Thomas KM, Prynn HW and Navarro J (1990) Molecular cloning of the Mel-Leu-Phe receptor from neurophilis (terramol) published in *J Biol Chem* 267:13780. *J Biol Chem* 265:20061–20064.
- Thomas KM, Taylor L and Navarro J (1991) The interleukin-8 receptor is encoded by a neurophilic-specific cDNA clone, FBR. *J Biol Chem* 266:14839–14841.
- Tiffany HL, Luens LA, Gao J-L, Pease J, Locati M, Comandini C, Modi W, Bonner TI and Murphy PM (1997) Identification of CCR2: A human monocyte and thymus receptor for the CC chemokine 1-509. *J Exp Med* 186:195–170.
- Topham PS, Ceizmadia V, Soler D, Hines D, Gerard CJ, Salant DJ and Hancock WW (1999) Lack of chemokine receptor CCR1 enhances Th1 responses and glomerular injury during nephrotic nephritis. *J Clin Invest* 104:1549–1557.
- Touramian S, Colin Y, Cuxton JP and Le Van Kim C (1995) Disruption of a GATA member in the Duffy gene promoter abolishes erythroid gene expression in Duffy-negative individuals. *Nat Genet* 10:224–228.
- Trentin L, Agostini C, Faccio M, Piazza F, Perin A, Bonvicini M, Gurrieri C, Galvan S, Adami F, Zambello R and Semenzato G (1999) The chemokine receptor CXCR3 is expressed on malignant B cells and mediates chemotaxis. *J Clin Invest* 104:115–121.
- Trkola A, Dragic T, Arthos J, Binley JM, Olson WC, Allaway GP, Cheng-Mayer C, Robinson J, Maddon PJ and Moore JP (1996) CD4-dependent, antibody-sensitive interactions between HIV-1 and its co-receptor CCR-5. *Nature (London)* 384:184–187.
- Tsao CL, Gludie RF, Carroll LA, Paradis T, Boyd JC, Nelson RT, Neote K and Charo IF (1998) Identification of C-C chemokine receptor 1 (CCR1) as the monocyte homologue of C-C chemokine (HCC)-1 receptor. *J Exp Med* 188:606–618.
- Ugueto EM, Hargrave M, Baldwin JM and Scherlier GF (1997) Arrangement of rhodopsin transmembrane alpha-helices. *Nature (London)* 389:203–206.
- Vanboute PM, Humphrey PPA and Spedding M (1998) NC-IUPHAR recommendations for nomenclature of receptors. In *The IUPHAR Compendium of Receptor Characterization and Classification*, 1st ed pp 31–33, Burlington Press, Foston, Colorado.
- Van Snick J, Housiaux F, Proost P, Van Damme J and Renauld JC (1996) 130e7 cell activation gene-3 chemokine protects murine T cell lymphomas against dexamethasone-induced apoptosis. *J Immunol* 157:2071–2076.
- Venema B, Zabolny A, Guetierres J, Garcia B, Roncal P, Albar JP, Ardrin C and Marquez G (1998) Molecular cloning, functional characterization and mRNA expression analysis of the murine chemokine receptor CCR6 and its specific ligand MIP-3alpha. *FEBS Lett* 440:188–194.

- Vieira J, Schall TJ, Corey L and Galabie AP (1998) Functional analysis of the human cytomegalovirus US28 gene by insertion mutagenesis with the green fluorescent protein. *J Virol* **72**:8158–8165.
- Virgin HW 4th, Latreille P, Wansley J, Hallsworth K, Weck KP, Dal Canto AJ and Speck SH (1997) Complete sequence and genomic analysis of murine gammaherpesvirus 68. *J Virol* **71**:5894–5904.
- Wakasugi K and Schimml P (1990) Two distinct cytokines released from a human antineoplastic cell line. *Science (Wash DC)* **248**:147–151.
- Walt A and Baggiolini M (1990) Generation of the neutrophil-activating peptide NAP-2 from platelet basic protein or connective tissue-activating peptide III through monocyte proteases. *J Exp Med* **171**:449–454.
- Walt A, Peverl F, Aschauer H and Baggiolini M (1987) Purification and amino acid sequencing of NAP, a new neutrophil-activating factor produced by monocytes. *Biochem Biophys Res Commun* **149**:755–761.
- Walt DA, Wu VV, de Lamo R, Dene H and McCoy LE (1977) Primary structure of human platelet factor 4. *Thromb Res **11**:859–858.*
- Wang JF, Liu ZY and Greenbaum D (1998) The alpha-chemokine receptor CXCR4 is expressed on the megakaryocytic lineage from progenitor to platelets and modulates migration and adhesion. *Blood* **92**:756–764.
- Weissman D, Rablin RL, Arthos J, Rubert A, Dybul M, Swofford R, Venkatesan S, Farber JM and Fauci AS (1997) Macrophage-tropic HIV and SIV envelope proteins induce a signal through the CCR5 chemokine receptor. *Nature (Lond)* **389**:981–985.
- Wells TN and Peitsch MC (1997) The chemokine information source: Identification and characterization of novel chemokines using the WorldWideWeb and expressed sequence tag databases. *J Leukoc Biol* **61**:545–550.
- Weng Y, Siciliano SJ, Waldburger KK, Sirotina-Melissar A, Staruch MJ, Daugherty BL, Gould SL, Springer MS and DeMartino JA (1998) Binding and functional properties of recombinant and endogenous CXCR3 chemokine receptors. *J Biol Chem* **273**:18288–18291.
- White JK, Shaw MA, Barton CH, Cerretti DP, Williams H, Mook BA, Carter NP, Peacock SC and Blackwell JM (1994) Genetic and physical mapping of 2436 in the region of the NRAMP and IL8R genes: Identification of a polymorphic repeat in exon 2 of NRAMP. *Genomics* **24**:352–362.
- White JR, Lee JM, Young BP, Hagerberg RP, Jurewicz AJ, Chaikina MA, Widdowson K, Foley JD, Martin LD, Griswold DE and Sarau HM (1998) Identification of a potent, selective non-peptide CXCR2 antagonist that inhibits interleukin-8-induced neutrophil migration. *J Biol Chem* **273**:10095–10098.
- Williamson K, Legler DF, Loescherer M, Ross RS, Delgado MB, Clark-Lewis I, Baggiolini M and Moser B (1998) The chemokine SLC is expressed in T cell areas of lymph nodes and mucosal lymphoid tissues and attracts activated T cells. *Eur J Cell Res* **18**:2025–2034.
- Winkler C, Modi W, Smith MW, Nelson GW, Wu X, Carrington M, Dean M, Hoque T, Tashiro K, Yabe D, Hughes MD, Vittinghoff II, Goedert JJ, O'Brien TR, Jacobson LP, Detels R, Donfield S, Willoughby A, Gomperts E, Vlahov D, Phair J, ALIVE Study, Hemophilia Growth and Development Study (HGDS), Multicenter AIDS Cohort Study (MACS), Multicenter Hemophilia Cohort Study (MCHS), San Francisco City Cohort (SFCC) and O'Brien SJ (1998) Genetic restriction of AIDS pathogenesis by an SDF-1 chemokine gene variant. *Science (Wash DC)* **279**:389–393.
- Wolpe SD and Cerami A (1989) Macrophage inflammatory proteins 1 and 2: Members of a novel superfamily of cytokines. *FASEB J* **3**:2656–2673.
- Wong LM, Myers SJ, Tsou CH, Gieding J, Arat H and Charo IF (1997) Organization and differential expression of the human monocyte chemoattractant protein 1 receptor gene: Evidence for the role of the carboxyl-terminal tail in receptor trafficking. *J Biol Chem* **272**:1038–1045.
- Wu L, LaRosa G and Mackay CR (1997) Interaction of chemokine receptor CXCR4 with its ligands: Multiple domains for HIV-1 gp120 binding and a single domain for chemokine binding. *J Exp Med* **186**:1373.
- Wuyts A, Van Osaeter L, Haelsa A, Samson I, Herdewijn P, Ben-Baruch A, Oppenheim JJ, Proest P and Van Damme J (1997) Characterization of synthetic human granulocyte chemotactic protein 2: Usage of chemokine receptors CXCR1 and CXCR2 and in vivo inflammatory properties. *Biochemistry* **36**:2716–2723.
- Xiao X, Wu L, Stanchove TS, Feng YR, Ugelstad S, Chen H, Shen Z, Riley JL, Broder CK, Sattentau QJ and Dimitrov DS (1998) Constitutive cell surface association between CD4 and CXCR4. *Proc Natl Acad Sci USA* **95**:7496–7501.
- Xu Y, Tanumura H, Arakaki R, Nakashima H, Zhang X, Fujii N, Uchiyama T and Hattori T (1998) Marked increase in anti-HIV entry mediated by CXCR4, linked to enhancement of the binding ability of tachyplesin analogs to CXCR4. *AIDS Res Hum Retrovir* **15**:19–27.
- Fangshars S, Komura E, Nagasawa J, Watarai H and Yamaguchi Y (1998) EBI1/CCR7 is a new member of dendritic cell chemokine receptor that is up-regulated upon maturation. *J Immunol* **161**:3096–3102.
- Yang AG, Bai X, Huang XF, Yao C and Chen S (1997) Phenotypic knockout of HIV type 1 chemokine receptor CCR5 by intrakines as potential therapeutic approach for HIV-1 infection. *Proc Natl Acad Sci USA* **94**:11567–11572.
- Yang D, Howard OM, Chen Q and Oppenheim JJ (1999a) Cutting edge: Immature dendritic cells generated from monocytes in the presence of TGF-beta 1 express functional C-C chemokine receptor 5. *J Immunol* **163**:737–741.
- Yang Y, Loy J, Ryscek RP, Carrasao D and Bravo R (1998) Antigen-induced eosinophilic lung inflammation depends on mice deficient in chemokine eotaxin. *Blood* **92**:3912–3923.
- Yang D, Chertov O, Ryckovaik SN, Chen Q, Buffo MD, Shogan J, Anderson M, Schroder JM, Wang JM, Howard OM and Oppenheim JJ (1999b) Beta-defensin: Linking innate and adaptive immunity through dendritic and T cell CCR6. *Science (Wash DC)* **286**:525–528.
- Yoneyama H, Harada A, Imai T, Baba M, Yoshie O, Zhang Y, Higashi H, Murai M, Asakura H and Matsushima K (1998) Pivotal role of TARC, a CC chemokine, in bacteria-induced inflammatory hepatic failure in mice. *J Clin Invest* **102**:1933–1941.
- Yoshida R, Imai T, Matsushima K, Yoshie O, Baba M, Kikura M, Nishimura M, Kikuchi M, Yamashita H and Yoshie O (1997) Molecular cloning of a novel human CC chemokine EBI1-lygand chemokine that is a specific functional ligand for EBI1. *CCRT, J Biol Chem* **272**:13803–13809.
- Yoshida R, Nagira M, Kikura M, Imai T, Matsushima K, Yoshie O (1998a) Second-ary lymphoid chemokine EBI1-lygand is a functional ligand for the CC chemokine receptor CCRT. *J Biol Chem* **273**:7118–7122.
- Yoshida T, Imai T, Kakizaki M, Nishimura M, Takagi S and Yoshie O (1998b) Identification of single C motif/lymphotactin receptor CXCR1. *J Biol Chem* **273**:16551–16554.
- Yoshida T, Imai T and Nomiyama H (1997) Novel lymphocyte-specific CC chemokines and their receptors. *J Leukoc Biol* **62**:634–644.
- Yoshimura T, Matsushima K, Tanaka S, Robinson EA, Appella E, Oppenheim JJ and Leonard EJ (1987) Purification of a human monocyte-derived neutrophil chemokine factor that has peptide sequence similarity to other host defense cytokines. *Nat Acad Sci USA* **84**:9233–9237.
- Yoshimura T, Yuhki N, Moore SC, Appella E, Lerman M and Leonard EJ (1989) Human monocyte chemoattractant protein-1 (MCP-1). Full-length cDNA cloning, expression in mitogen-stimulated blood mononuclear leukocytes, and sequence similarity to mouse chemokine gene JE. *FEBS Lett* **244**:487–495.
- Youn BS, Kim CH, Smith FO and Broxmeyer HE (1999) TECK, an efficacious chemoattractant for human thymocytes, uses GPR9–6/CCR5 as a specific receptor. *Blood* **94**:2533–2536.
- Youn BS, Zhang SM, Lee EK, Park DH, Broxmeyer HF, Murphy PM, Locati M, Pease JE, Kim KJ, Kim K and Kwon BS (1997) Molecular cloning of leukotactin-1, a novel human beta-chemokine, a chemoattractant for neutrophils, monocytes, and lymphocytes, and a potent agonist at CC chemokine receptors 1 and 3. *J Immunol* **159**:5201–5205.
- Yoshida R, Imai T, Matsushima K, Golding H and Farber JM (2000) CCR5 and CCR9: Two receptors for the chemokine CCL25/TECK/Ck1b-1 that differ in their sensitivity to ligand. *J Immunol* **164**:1293–1305.
- Zaballo A, Gutierrez J, Varona R, Arduini C and Marquez G (1999) Cutting edge: Identification of the orphan chemokine receptor GPR-96 as CCR9, the receptor for the chemokine TECK. *J Immunol* **163**:5971–5975.
- Zabel BA, Agace WM, Campbell JJ, Heath HM, Parent D, Roberts AL, Ebert EC, Kassam N, Qin S, Zovko M, LaRosa GJ, Yang LL, Solor D, Butcher EC, Ponath PD, Parker CM and Andrew DP (1999) Human G protein-coupled receptor GPR-96/CCR chemokine receptor 9 is selectively expressed on intestinal homing T lymphocytes, mucosal lymphocytes, and T-cells and is required for thymus-expressed chemokine-mediated chemotaxis. *J Exp Med* **190**:1241–1256.
- Zaitseva M, Blauvelt A, Lee S, Lapham CK, Klaus-Kortum V, Mostowski H, Maniashewitz J and Golding H (1997) Expression and function of CCR5 and CXCR4 on human Langerhans cells and macrophages: Implications for HIV primary infection. *Nat Med* **3**:1369–1376.
- Zaitseva MB, Lee S, Rabini RL, Tiffany HL, Farber JM, Peden KW, Murphy PM and Golding H (1998) CXCR4 and CCR5 on human thymocytes: Biological function and role in HIV-1 infection. *J Immunol* **161**:3103–3113.
- Zhang JY, Dragic T, Cao Y, Kostrikis L, Kwon DS, Littman DR, Kowal Raman VN and Moore JP (1998a) Use of coreceptors other than CCR5 by non-syncretism-inducing adult and pediatric isolates of human immunodeficiency virus type 1 are rare in vitro. *J Virol* **72**:9337–9344.
- Zhang L, He T, Talal A, Wang G, Frankel SS and Ho DD (1998b) In vivo distribution of the human immunodeficiency virus/simian immunodeficiency virus coreceptors: CXCR4, CCR5, and CCR2. *J Virol* **72**:5035–5045.
- Zhang S, Youn BS, Gao JJ, Murphy PM and Kwon BS (1999) Differential effects of leukotactin-1 and chemokine inflammatory protein-1 alpha on neutrophils mediated by CCR1. *J Immunol* **163**:4938–4942.
- Zhou Y, Kurahara T, Ryscek RP, Yang Y, Ryan C, Loy J, Warg G and Bravo R (1998) Impaired macrophage function and enhanced T cell-dependent immune response in mice lacking CCR5: The mouse homologue of the major HIV-1 coreceptor. *J Immunol* **160**:4018–4025.
- Zimmerman PA, Buckner-White A, Alkhatib G, Spalding T, Kubacki J, Combadiere C, Weissman D, Cohen O, Rubbert A, Lam G, Vaccarezza M, Kennedy PE, Kumararajam V, Giorgi JV, Detels R, Hunter J, Chopel M, Berger EA, Fauci AS, Nutman TB and Murphy PM (1997) Inherited resistance to HIV-1 conferred by an inactivating mutation in CC chemokine receptor 5: Studies in populations with contrasting clinical phenotypes, defined racial background and quantified risk. *Mol Med* **3**:23–36.
- Zimmerman PA, Woolley I, Masidine GL, Miller SM, McNamara DT, Hazlett F, Mgone SC, Alpers MP, Genton B, Boatin BA and Kaurara JW (1999) Emergence of FV*Anal in a *Plasmodium vivax*-endemic region of Papua New Guinea. *Proc Natl Acad Sci USA*, in press.
- Zingoni A, Soto H, Hedrick JA, Stoppaccaro A, Storazzi CT, Sinigaglia F, D'Ambrosio D, O'Garra A, Robinson D, Rocchi M, Santoni A, Zlotnik A and Napolitano M (1998) The chemokine receptor CCR6 is preferentially expressed in Th2 but not Th1 cells. *J Immunol* **161**:547–551.
- Zlotnik A, Morales J and Hedrick JA (1999) Recent advances in chemokines and chemokine receptors. *Crit Rev Immunol* **19**:1–47.
- Zou P, Isewawa Y, Nakano K, Haque M, Horiguchi Y and Yamanishi K (1999) Human herpesvirus 6 open reading frame US3 encodes a functional chemokine. *J Virol* **73**:1000–1004.
- Zou YR, Kottmann AH and Littman DR (1998) Function of the chemokine receptor CXCR4 in hematopoiesis and in cerebellar development. *Nature (Lond)* **393**:595–598.